



## Research Article

# Fractional derivative analysis of Asthma with the effect of environmental factors

Rashid JAN<sup>1</sup>, M. S. ZOBAER<sup>2</sup>, Şuayip YÜZBAŞI<sup>3</sup>, Attaullah<sup>4</sup>, Muhammad JAWAD<sup>1</sup>, Asif JAN<sup>5</sup>

<sup>1</sup>Department of Mathematics, University of Swabi, Swabi 23561, Pakistan

<sup>2</sup>McGovern Medical School, The University of Texas Health Science Center at Houston, Texas 77030, USA

<sup>3</sup>Department of Mathematics, Faculty of Science, Bartın University Tr-74100 Bartın, Türkiye

<sup>4</sup>Department of Mathematics, University of Chitral, Chitral, 17200, KP, Pakistan

<sup>5</sup>Department of Pathogenic Microbiology & Immunology, School of Basic medical Sciences, Xi'an

## ARTICLE INFO

### Article history

Received: 30 June 2021

Revised: 11 August 2021

Accepted: 18 September 2021

### Keywords:

Asthma Disease; Environmental Factors; Mathematical Model; Numerical Method; Dynamical Behaviour

## ABSTRACT

It is observed that the exposure to environmental factors such as indoors and outdoors air pollution, cigarette smoke, and allergens are highly related to asthma attacks. It is also reported that limited exposure to asthma triggers, cure due to medicine, the attacks of asthma can be minimized. In this paper, we formulate the dynamics of asthma with smoking and environmental factors classes in the fractional Caputo-Fabrizio (CF) framework to visualize its dynamical behaviour. We delineate the important properties of the CF derivative for the analysis of our model. The model is then analyzed for the basic properties and the uniqueness and existence of the hypothesized asthma system are investigated via the theory of fixed point. Furthermore, a novel numerical scheme is presented for the solution of our fractional system to illustrate the time series of asthma model. The dynamical behaviour of our asthma model is then highlighted numerically to show the impact of fractional-order  $\vartheta$  on the system and to visualize the role of input factors on the dynamics of asthma disease.

**Cite this article as:** Jan R, Zobaer MS, Yüzbaşı Ş, Attaullah, Jawad M, Jan A. Fractional derivative analysis of Asthma with the effect of environmental factors. Sigma J Eng Nat Sci 2024;42(1):177–188.

## INTRODUCTION

It is reported that asthma is a chronic disease that causes the inside walls of the airways of humans and make it hard to breathe. This chronic disease makes it hard to breathe and shorten the breath when the infected individual breathes out. In some cases asthma is a mild annoyance while it may be a big problem for others that disturb their

daily life and is life threatening in case of asthma attack. There is no specific treatment for asthma, however, its signs can be monitored. The patients with asthma should consult with the doctors and can modify care according to the condition. The symptoms and signs of asthma are chest tightness or pain, wheezing are coughing attacks, such as flu and cold, infect respiratory system, shortness of breath,

### \*Corresponding author.


\*E-mail address: [suayipyuzbasi@gmail.com](mailto:suayipyuzbasi@gmail.com)

This paper was recommended for publication in revised form by Regional Editor Vildan Çetkin



sleeping problems related to shortness of breath, wheezing or coughing. Furthermore, wheezing in children is a common sign of asthma during exhaling, and increasing difficulty breathing. The lack of oxygen in asthma asthma make the patient serious and can feel chest pain or even loss of consciousness. In Figure 1, a systematic flow chart of asthma is given to represent the phenomena of asthma from trigger factors to signs and symptoms of asthma.

It is reported that asthma can be induced not only by air contaminants such as di-isocyanates, hexachloroplatinates, etc., but cigarette smoke also triggers asthma [1-3]. Our body can be affected in many ways by smoking which is dangerous to the lungs. This contributes to reduced lung function that makes the lungs vulnerable to asthma triggers. High concentrations of irritants including nitrogen oxides, ammonia, acrolein and formaldehyde are present in cigarette smoke. The expert insisted that lungs damage occurs not only from mainstream smoke but also from side stream smoke which comes to the host through a nearby smoker in the society. Therefore, these unpleasant substances can set off an asthma attack when the smoke of tobacco inhaled by a person, whether for personal smoking or passive smoke. More than 15 million children per day are threatened and open to smoke, and more than 1,000,000 children with asthma become more serious after second-hand smoke exposure [4-5]. Also in the cause of asthma, contaminants released in the air such as vanadium and nickel particles from power stations and oil refineries, heavy metal cocktails from SLF burning cement, brick and lime works and dust from open cast coal mining and coal fired cement and other works play important roles [6-8]. Thus, a prone population, due to constant exposure to air contaminants, they become asthmatic. Mathematical models play an important role to explore the dynamics of diseases [9-13, 50-51] and produce significant information about the dynamics of disease for control and prevention [14-17]. Different numerical approaches has been developed by researchers



**Figure 1.** Illustration of systematic flow of asthma disease from trigger factors to signs and symptoms.

to investigate the dynamics of diseases [18-22, 51]. There is limited literature on modeling the dynamics of asthma to visualize, conceptualize and understand the transmission phenomena of asthma [23-27]. The dynamics of asthma has been recently investigated by the researchers in their work from different aspects [28-29]. It is noted that in the development and spread of asthma, continuous exposure to smoking plays a very important role. This very significant factor has been accounted for in our model by considering a different class of smokers.

It is crystal clear that fractional systems give knowledge about the dynamics of various diseases more accurate, broader, precious and precise than integer systems [30,31]. It surpasses integer-order models due to hereditary qualities and memory definition [35-36], moreover, fractional models better explore and demonstrate the dynamics between two points. In the literature, it has been demonstrated that fractional models provide the best match to real data [37,38]. These new theories have been efficiently used in physics, economics, biology, engineering and many other fields to model natural processes [39-58]. These operators, on the other hand, have a power law kernel and are limited in their ability to mimic physical situations. Caputo and Fabrizio recently proposed an additional fractional differential operator with an exponential decay kernel to address this challenge. The Caputo-Fabrizio (CF) operator is a revolutionary technique to fractional derivative that has drawn the attention of numerous researchers owing to its non-singular kernel. In addition, the CF operator is ideal for simulating particular types of real-world problems that follow the exponential decay law. Developing a mathematical model employing the CF fractional order derivative has become a fascinating subject of the upcoming research.

We choose to use fractional derivative to investigate the dynamics of asthma with smoking and environmental class, motivated by the accurate results of the fractional derivative. We represent the fundamental results and theory of fractional CF derivative in 1 Section of the article. In the second section, using the CF derivative, a fractional model for asthma infection with smoking and environmental factor compartment is developed. Mathematical skills are then used to investigate the hypothesised asthma model. In Section 3, the uniqueness and existence of the solution of the formulated FO model of asthma through the fixed-point theorem is presented. The dynamics of proposed asthma model is then investigated with the effect of the different input parameter numerically in Section 4. Finally, in the last section, concluding observations and ideas are offered.

## FORMULATION OF THE ASTHMA MODEL

In the evaluation of the dynamics, the total population size  $N(t)$  is divided into four subclasses susceptibles class  $\mathcal{A}_S(t)$ , exposed class  $\mathcal{A}_E(t)$ , asthma infected class  $\mathcal{A}_I(t)$  and smokers class  $\mathcal{A}_C(t)$ , while the cumulative concentration

of pollutants present in the environment is indicated by  $\mathcal{A}_p(t)$ . The recruitment rate of susceptible and smoker is indicated by  $\Pi$  and  $\Xi$ , respectively while the natural death rate is assumed to be  $\mu$  for all classes of humans population. We supposed that the susceptible population become asthmatic by continuous exposure to air pollutants such as diisocyanates, hexachloroplatinates, etc. The susceptible humans population become exposed at a rate  $\beta$  due to smoking while a fraction  $\gamma$  of susceptible moves to the exposed class due to environmental factors. In the exposed class, two types of transmission has been observed denoted by  $\lambda_1$  and  $\lambda_2$ .  $\lambda_1$  is the transmission rate to the infected group from the exposed group due to smokers while  $\lambda_2$  is the transmission rate from the infected group to the exposed group due to environmental factors. Moreover, we assume  $\eta_1$  be the rate at which smokers become infected while  $\eta_2$  is the rate at which smokers become infected through environmental factors. We indicated the asthma infection induced death rate by  $\alpha$  in our formulation. We symbolize the emission rate of pollutants by  $A(N)$  which depend on population, however, we consider it to be constant. The rate of natural depletion is denoted by  $\nu$  while smokers emit fumes rate is given by  $q$  and the rate at which smoker leave smoking is  $\eta$ . The following system of ODE's describes the dynamics of asthma

$$\begin{cases} \frac{d\mathcal{A}_S}{dt} = \Pi - \beta\mathcal{A}_S\mathcal{A}_C - \gamma\mathcal{A}_S\mathcal{A}_P - \mu\mathcal{A}_S, \\ \frac{d\mathcal{A}_E}{dt} = \beta\mathcal{A}_S\mathcal{A}_C + \gamma\mathcal{A}_S\mathcal{A}_P - \lambda_1\mathcal{A}_E\mathcal{A}_P - \lambda_2\mathcal{A}_E\mathcal{A}_C - \mu\mathcal{A}_E, \\ \frac{d\mathcal{A}_I}{dt} = \lambda_1\mathcal{A}_E\mathcal{A}_P + \lambda_2\mathcal{A}_E\mathcal{A}_C + \eta_1\mathcal{A}_C + \eta_2\mathcal{A}_C\mathcal{A}_P - \alpha\mathcal{A}_I - \mu\mathcal{A}_I, \\ \frac{d\mathcal{A}_C}{dt} = \Xi - \eta_1\mathcal{A}_C - \eta_2\mathcal{A}_C\mathcal{A}_P - \eta\mathcal{A}_C - \mu\mathcal{A}_C, \\ \frac{d\mathcal{A}_P}{dt} = A(N) - \nu\mathcal{A}_P + q\mathcal{A}_C, \end{cases} \quad (1)$$

with appropriate initial condition for vector

$$\mathcal{A}_S(0) \geq 0, \mathcal{A}_E(0) \geq 0, \mathcal{A}_I(0) \geq 0, \mathcal{A}_C(0) \geq 0, \mathcal{A}_P(0) \geq 0. \quad (2)$$

It is cristal clear that fractional order systems provide more dependable, deeper, valuable, and accurate data concerning a disease's dynamics. To be more specific, the effect of memory can be captured through fractional operators which is mostly occurs in biological phenomena. Moreover, fractional order parameter make it more flexible for data fitting. Therefore, the above system (1) with (2) of asthma disease is represented in the framework of CF derivative in the following manner

$$\begin{aligned} {}^{\text{CF}}_0D_t^\varsigma \mathcal{A}_S &= \Pi - \beta\mathcal{A}_S\mathcal{A}_C - \gamma\mathcal{A}_S\mathcal{A}_P - \mu\mathcal{A}_S; \\ {}^{\text{CF}}_0D_t^\varsigma \mathcal{A}_E &= \beta\mathcal{A}_S\mathcal{A}_C + \gamma\mathcal{A}_S\mathcal{A}_P - \lambda_1\mathcal{A}_E\mathcal{A}_P - \lambda_2\mathcal{A}_E\mathcal{A}_C - \mu\mathcal{A}_E; \\ {}^{\text{CF}}_0D_t^\varsigma \mathcal{A}_I &= \lambda_1\mathcal{A}_E\mathcal{A}_P + \lambda_2\mathcal{A}_E\mathcal{A}_C + \eta_1\mathcal{A}_C + \eta_2\mathcal{A}_C\mathcal{A}_P - \alpha\mathcal{A}_I - \mu\mathcal{A}_I; \end{aligned}$$

$$\begin{aligned} {}^{\text{CF}}_0D_t^\varsigma \mathcal{A}_C &= \Xi - \eta_1\mathcal{A}_C - \eta_2\mathcal{A}_C\mathcal{A}_P - \eta\mathcal{A}_C - \mu\mathcal{A}_C; \\ {}^{\text{CF}}_0D_t^\varsigma \mathcal{A}_P &= A(N) - \nu\mathcal{A}_P + q\mathcal{A}_C; \end{aligned} \quad (3)$$

where  $\vartheta$  is the order of CF fractional derivative such that  $0 < \vartheta \leq 1$ . In the upcoming subsection of the article, we represent basic idea and concept of fractional derivative theory that will be utilized in the analysis of our proposed fractional-order model of asthma infection.

### Basic Fractional Theory

In this subsection of the article, we will list fundamental results and theory of CF fractional derivative for the analysis of our asthma disease model with environmental factors. We take start from the following theorem:

**Definition 1** Assume a function  $g$  such that  $g \in H^1(a,b)$  with the condition that  $a < b$ , then the fractional derivative in CF framework [48] is given by

$$D_t^\varsigma(g(t)) = \frac{V(\varsigma)}{1-\varsigma} \int_a^t g'(x) \exp[-\varsigma \frac{t-x}{1-\varsigma}] dx, \quad (4)$$

with fractional-order  $\varsigma$  with the condition that  $\varsigma \in [0,1]$  and  $V(\tau)$  indicates normality such that  $V(0) = V(1) = 1$  [48]. Here, if  $g \notin H^1(a,b)$ , then we have the below fractional form

$$D_t^\varsigma(g(t)) = \frac{\varsigma V(\varsigma)}{1-\varsigma} \int_a^t (g(t) - g(x)) \exp[-\varsigma \frac{t-x}{1-\varsigma}] dx. \quad (5)$$

**Remark 1** Assume the value  $\alpha = \frac{1-\varsigma}{\varsigma} \in [0, \infty)$  and  $\varsigma = \frac{1}{1+\alpha} \in [0,1]$ , then the above (5) takes the below form

$$D_t^\varsigma(g(t)) = \frac{M(\alpha)}{\alpha} \int_a^t g'(x) e^{[-\frac{t-x}{\alpha}]} dx, \quad (6)$$

in which  $M(0) = M(\infty) = 1$ . Furthermore, we have the below

$$\lim_{\alpha \rightarrow 0} \frac{1}{\alpha} \exp[-\frac{t-x}{\alpha}] = \delta(x - t).$$

Here, the definition of fractional integral for CF framework will be represented see in [48].

**Definition 2** Assume a function  $g$ , and  $\varsigma$  be the fractional-order then the fractional integral is

$$I_t^\varsigma(g(t)) = \frac{2(1-\varsigma)}{(2-\varsigma)V(\varsigma)} g(t) + \frac{2\varsigma}{(2-\varsigma)V(\varsigma)} \int_0^t g(u) du, \quad t \geq 0. \quad (7)$$

with the condition  $0 < \varsigma < 1$  [49].

**Remark 2** Here, we investigate the above mentioned 2 and get the following

$$\frac{2(1-\varsigma)}{(2-\varsigma)V(\varsigma)} + \frac{2\varsigma}{(2-\varsigma)V(\varsigma)} = 1, \quad (8)$$

which further implies that  $V(\zeta) = \frac{2}{2-\zeta}$ ,  $0 < \zeta < 1$ . In the upcoming step, we represent a novel fractional derivative introduced in [49] by using (8) and is as

$$D_t^\zeta(g(t)) = \frac{1}{1-\zeta} \int_0^t g'(x) \exp[\zeta \frac{t-x}{1-\zeta}] dx, 0 < \zeta < 1.$$

**NUMERICAL SCHEME AND ANALYSIS**

Here, the solution of the proposed asthma disease model will be investigated for existence through fixed-point theory. We use the concept of CF fractional derivative on the system (3), and get the following

$$\begin{aligned} A_S(t) - A_S(0) &= {}_0^{\text{CF}}I_t^\zeta [\Pi - \beta A_S A_C - \gamma A_S A_P - \mu A_S], \\ A_E(t) - A_E(0) &= {}_0^{\text{CF}}I_t^\zeta [\beta A_S A_C + \gamma A_S A_P - \lambda_1 A_E A_P - \lambda_2 A_E A_C - \mu A_E], \\ A_I(t) - A_I(0) &= {}_0^{\text{CF}}I_t^\zeta [\lambda_1 A_E A_P + \lambda_2 A_E A_C + \eta_1 A_C + \eta_2 A_C A_P - \alpha A_I - \mu A_I], \\ A_P(t) - A_P(0) &= {}_0^{\text{CF}}I_t^\zeta [\Xi - \eta_1 A_C - \eta_2 A_C A_P - \eta A_C - \mu A_C], \\ A_P(t) - A_P(0) &= {}_0^{\text{CF}}I_t^\zeta [A(N) - \nu A_P + q A_C]. \end{aligned} \tag{9}$$

Apply the idea presented in [49], the above (9) implies

$$\begin{aligned} A_S(t) - A_S(0) &= \frac{2(1-\theta)}{(2-\theta)U(\theta)} [\Pi - \beta A_S A_C - \gamma A_S A_P - \mu A_S] \\ &\quad + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t [\Pi - \beta A_S A_C - \gamma A_S A_P - \mu A_S] dy, \\ A_E(t) - A_E(0) &= \frac{2(1-\theta)}{(2-\theta)U(\theta)} [\beta A_S A_C + \gamma A_S A_P - \lambda_1 A_E A_P - \lambda_2 A_E A_C - \mu A_E] \\ &\quad + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t [\beta A_S A_C + \gamma A_S A_P - \lambda_1 A_E A_P - \lambda_2 A_E A_C - \mu A_E] dy, \\ A_I(t) - A_I(0) &= \frac{2(1-\theta)}{(2-\theta)U(\theta)} [\lambda_1 A_E A_P + \lambda_2 A_E A_C + \eta_1 A_C + \eta_2 A_C A_P - \alpha A_I - \mu A_I] \\ &\quad + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t [\lambda_1 A_E A_P + \lambda_2 A_E A_C + \eta_1 A_C + \eta_2 A_C A_P - \alpha A_I - \mu A_I] dy, \\ A_C(t) - A_C(0) &= \frac{2(1-\theta)}{(2-\theta)U(\theta)} [\Xi - \eta_1 A_C - \eta_2 A_C A_P - \eta A_C - \mu A_C] \\ &\quad + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t [\Xi - \eta_1 A_C - \eta_2 A_C A_P - \eta A_C - \mu A_C] dy, \\ A_P(t) - A_P(0) &= \frac{2(1-\theta)}{(2-\theta)U(\theta)} [A(N) - \nu A_P + q A_C] + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t [A(N) - \nu A_P + q A_C] dy. \end{aligned} \tag{10}$$

In the next step, we proceed in the following manner

$$\begin{cases} \mathcal{L}_1(t, \mathcal{A}_S) &= \Pi - \beta \mathcal{A}_S \mathcal{A}_C - \gamma \mathcal{A}_S \mathcal{A}_P - \mu \mathcal{A}_S, \\ \mathcal{L}_2(t, \mathcal{A}_E) &= \beta \mathcal{A}_S \mathcal{A}_C + \gamma \mathcal{A}_S \mathcal{A}_P - \lambda_1 \mathcal{A}_E \mathcal{A}_P - \lambda_2 \mathcal{A}_E \mathcal{A}_C - \mu \mathcal{A}_E, \\ \mathcal{L}_3(t, \mathcal{A}_I) &= \lambda_1 \mathcal{A}_E \mathcal{A}_P + \lambda_2 \mathcal{A}_E \mathcal{A}_C + \eta_1 \mathcal{A}_C + \eta_2 \mathcal{A}_C \mathcal{A}_P - \alpha \mathcal{A}_I - \mu \mathcal{A}_I, \\ \mathcal{L}_4(t, \mathcal{A}_C) &= \Xi - \eta_1 \mathcal{A}_C - \eta_2 \mathcal{A}_C \mathcal{A}_P - \eta \mathcal{A}_C - \mu \mathcal{A}_C, \\ \mathcal{L}_5(t, \mathcal{A}_P) &= A(N) - \nu \mathcal{A}_P + q \mathcal{A}_C, \end{cases}$$

**Theorem 1** *If the following condition satisfies*

$$0 \leq (\beta M + \gamma N + \mu) < 1,$$

then the kernels  $\mathcal{L}_1, \mathcal{L}_2, \mathcal{L}_3, \mathcal{L}_4$  and  $\mathcal{L}_5$  assures the Lipschitz condition.

**Proof.** For the proof of Theorem 2, we first take  $\mathcal{A}_S$  and  $\mathcal{A}_{S1}$ , and start from  $\mathcal{L}_1$  in the following manner

$$\begin{aligned} \mathcal{L}_1(t, \mathcal{A}_S) - \mathcal{L}_1(t, \mathcal{A}_{S1}) &= -\beta \mathcal{A}_S \mathcal{A}_C - \gamma \mathcal{A}_S \mathcal{A}_P - \mu \mathcal{A}_S \\ &\quad + \beta \mathcal{A}_{S1} \mathcal{A}_C + \gamma \mathcal{A}_{S1} \mathcal{A}_P + \mu \mathcal{A}_{S1}, \end{aligned}$$

this implies that

$$\begin{aligned} \mathcal{L}_1(t, \mathcal{A}_S) - \mathcal{L}_1(t, \mathcal{A}_{S1}) &= \beta(\mathcal{A}_{S1} - \mathcal{A}_S) \mathcal{A}_C + \gamma(\mathcal{A}_{S1} - \mathcal{A}_S) \mathcal{A}_P \\ &\quad + \mu(\mathcal{A}_{S1} - \mathcal{A}_S) \end{aligned} \tag{11}$$

Here, apply norm on Eq. (11) and simplify, we get the below

$$\begin{aligned} \|\mathcal{L}_1(t, \mathcal{A}_S) - \mathcal{L}_1(t, \mathcal{A}_{S1})\| &\leq \|\beta(\mathcal{A}_{S1} - \mathcal{A}_S) \mathcal{A}_C\| + \|\gamma(\mathcal{A}_{S1} - \mathcal{A}_S) \mathcal{A}_P\| \\ &\quad + \|\mu(\mathcal{A}_{S1} - \mathcal{A}_S)\|, \leq \beta \|\mathcal{A}_{S1} - \mathcal{A}_S\| \\ &\quad \|\mathcal{A}_C\| + \gamma \|\mathcal{A}_{S1} - \mathcal{A}_S\| \|\mathcal{A}_P\| + \mu \|\mathcal{A}_{S1} - \mathcal{A}_S\|, \\ &\leq \beta M \|\mathcal{A}_{S1} - \mathcal{A}_S\| + \gamma N \\ &\quad \|\mathcal{A}_{S1} - \mathcal{A}_S\| + \mu \|\mathcal{A}_{S1} - \mathcal{A}_S\|, \\ &\leq (\beta M + \gamma N + \mu) \|\mathcal{A}_{S1} - \mathcal{A}_S\|. \end{aligned}$$

At this stage, we assume  $\mu_1 = (\beta M + \gamma N + \mu)$ , then the following result is obtained

$$\|\mathcal{L}_1(t, \mathcal{A}_S) - \mathcal{L}_1(t, \mathcal{A}_{S1})\| \leq \mu_1 \|\mathcal{A}_S(t) - \mathcal{A}_S(t_1)\|.$$

Thus the Lipschitz condition is fulfilled for  $\mathcal{L}_1$ , in addition to this the condition  $0 \leq (\beta M + \gamma N + \mu) < 1$  assure the contraction is also satisfied. In the same way, the Lipschitz conditions for the other cases of our system are determined as

$$\begin{aligned} \|\mathcal{L}_2(t, \mathcal{A}_E) - \mathcal{L}_2(t, \mathcal{A}_{E1})\| &\leq \mu_2 \|\mathcal{A}_E(t) - \mathcal{A}_E(t_1)\|, \\ \|\mathcal{L}_3(t, \mathcal{A}_I) - \mathcal{L}_3(t, \mathcal{A}_{I1})\| &\leq \mu_3 \|\mathcal{A}_I(t) - \mathcal{A}_I(t_1)\|, \\ \|\mathcal{L}_4(t, \mathcal{A}_C) - \mathcal{L}_4(t, \mathcal{A}_{C1})\| &\leq \mu_4 \|\mathcal{A}_C(t) - \mathcal{A}_C(t_1)\|, \\ \|\mathcal{L}_5(t, \mathcal{A}_P) - \mathcal{L}_5(t, \mathcal{A}_{P1})\| &\leq \mu_5 \|\mathcal{A}_P(t) - \mathcal{A}_P(t_1)\|. \end{aligned} \tag{12}$$

Further simplification of Eq. (10) implies

$$\begin{cases} \mathcal{A}_S(t) &= \mathcal{A}_S(t)(0) + \frac{2(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_1(t, \mathcal{A}_S(t)) + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_1(y, \mathcal{A}_S(y))) dy, \\ \mathcal{A}_E(t) &= \mathcal{A}_E(t)(0) + \frac{2(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_2(t, \mathcal{A}_E(t)) + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_2(y, \mathcal{A}_E(y))) dy, \\ \mathcal{A}_I(t) &= \mathcal{A}_I(t)(0) + \frac{2(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_3(t, \mathcal{A}_I(t)) + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_3(y, \mathcal{A}_I(y))) dy, \\ \mathcal{A}_C(t) &= \mathcal{A}_C(t)(0) + \frac{2(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_4(t, \mathcal{A}_C(t)) + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_4(y, \mathcal{A}_C(y))) dy, \\ \mathcal{A}_P(t) &= \mathcal{A}_P(t)(0) + \frac{2(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_5(t, \mathcal{A}_P(t)) + \frac{2\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_5(y, \mathcal{A}_P(y))) dy, \end{cases} \tag{13}$$

its recursive form is given by

$$\begin{cases} \mathcal{A}_{Sn}(t) &= 2 \frac{(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_1(t, \mathcal{A}_{S(n-1)}) + 2 \frac{\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_1(y, \mathcal{A}_{S(n-1)})) dy, \\ \mathcal{A}_{En}(t) &= 2 \frac{(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_2(t, \mathcal{A}_{E(n-1)}) + 2 \frac{\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_2(y, \mathcal{A}_{E(n-1)})) dy, \\ \mathcal{A}_{In}(t) &= 2 \frac{(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_3(t, \mathcal{A}_{I(n-1)}) + 2 \frac{\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_3(y, \mathcal{A}_{I(n-1)})) dy, \\ \mathcal{A}_{Cn}(t) &= 2 \frac{(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_4(t, \mathcal{A}_{C(n-1)}) + 2 \frac{\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_4(y, \mathcal{A}_{C(n-1)})) dy, \\ \mathcal{A}_{Pn}(t) &= 2 \frac{(1-\theta)}{(2-\theta)U(\theta)} \mathcal{L}_5(t, \mathcal{A}_{P(n-1)}) + 2 \frac{\theta}{(2-\theta)U(\theta)} \int_0^t (\mathcal{L}_5(y, \mathcal{A}_{P(n-1)})) dy, \end{cases} \tag{14}$$

with proper initial conditions

$$\begin{aligned} \mathcal{A}_S^0(t) &= \mathcal{A}_S(0), \mathcal{A}_E^0(t) = \mathcal{A}_E(0), \mathcal{A}_I^0(t) = \mathcal{A}_I(0), \\ \mathcal{A}_C^0(t) &= \mathcal{A}_C(0), \mathcal{A}_P^0(t) = \mathcal{A}_P(0). \end{aligned}$$

Here, the succeeding difference form is

$$\begin{aligned}
 \hat{h}_{1n}(t) &= \mathcal{A}_{S_n}(t) - \mathcal{A}_{S_{(n-1)}}(t) \\
 &= \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_1(t, \mathcal{A}_{S_{(n-1)}}) - \mathcal{L}_1(t, \mathcal{A}_{S_{(n-2)}})) \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \int_0^t (\mathcal{L}_1(y, \mathcal{A}_{S_{(n-1)}}) - \mathcal{L}_1(y, \mathcal{A}_{S_{(n-2)}})) dy, \\
 \hat{h}_{2n}(t) &= \mathcal{A}_{E_n}(t) - \mathcal{A}_{E_{(n-1)}}(t) \\
 &= \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_2(t, \mathcal{A}_{E_{(n-1)}}) - \mathcal{L}_2(t, \mathcal{A}_{E_{(n-2)}})) \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \int_0^t (\mathcal{L}_2(y, \mathcal{A}_{E_{(n-1)}}) - \mathcal{L}_2(y, \mathcal{A}_{E_{(n-2)}})) dy, \\
 \hat{h}_{3n}(t) &= \mathcal{A}_{I_n}(t) - \mathcal{A}_{I_{(n-1)}}(t) \\
 &= \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_3(t, \mathcal{A}_{I_{(n-1)}}) - \mathcal{L}_3(t, \mathcal{A}_{I_{(n-2)}})) \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \int_0^t (\mathcal{L}_3(y, \mathcal{A}_{I_{(n-1)}}) - \mathcal{L}_3(y, \mathcal{A}_{I_{(n-2)}})) dy, \\
 \hat{h}_{4n}(t) &= \mathcal{A}_{C_n}(t) - \mathcal{A}_{C_{(n-1)}}(t) \\
 &= \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_4(t, \mathcal{A}_{C_{(n-1)}}) - \mathcal{L}_4(t, \mathcal{A}_{C_{(n-2)}})) \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \int_0^t (\mathcal{L}_4(y, \mathcal{A}_{C_{(n-1)}}) - \mathcal{L}_4(y, \mathcal{A}_{C_{(n-2)}})) dy, \\
 \hat{h}_{5n}(t) &= \mathcal{A}_{P_n}(t) - \mathcal{A}_{P_{(n-1)}}(t) \\
 &= \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_5(t, \mathcal{A}_{P_{(n-1)}}) - \mathcal{L}_5(t, \mathcal{A}_{P_{(n-2)}})) \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \int_0^t (\mathcal{L}_5(y, \mathcal{A}_{P_{(n-1)}}) - \mathcal{L}_5(y, \mathcal{A}_{P_{(n-2)}})) dy.
 \end{aligned} \tag{15}$$

Here, we observe the following

$$\begin{cases}
 \mathcal{A}_{S_n}(t) = \sum_{i=1}^n \hat{h}_{1i}(t), \mathcal{A}_{E_n}(t) = \sum_{i=1}^n \hat{h}_{2i}(t), \mathcal{A}_{I_n}(t) = \sum_{i=1}^n \hat{h}_{3i}(t), \\
 \mathcal{A}_{C_n}(t) = \sum_{i=1}^n \hat{h}_{4i}(t), \mathcal{A}_{P_n}(t) = \sum_{i=1}^n \hat{h}_{5i}(t).
 \end{cases}$$

Following the same way, we have

$$\begin{aligned}
 \|\hat{h}_{1n}(t)\| &= \|\mathcal{A}_{S_n}(t) - \mathcal{A}_{S_{(n-1)}}(t)\| = \left\| 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_1(t, \mathcal{A}_{S_{(n-1)}}) \right. \\
 &\quad \left. - \mathcal{L}_1(t, \mathcal{A}_{S_{(n-2)}})) + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \int_0^t (\mathcal{L}_1(y, \mathcal{A}_{S_{(n-1)}}) \right. \\
 &\quad \left. - \mathcal{L}_1(y, \mathcal{A}_{S_{(n-2)}})) dy \right\|.
 \end{aligned} \tag{16}$$

By triangular inequality, Eq. (16) becomes

$$\begin{aligned}
 \|\mathcal{A}_{S_n}(t) - \mathcal{A}_{S_{(n-1)}}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \|\mathcal{L}_1(t, \mathcal{A}_{S_{(n-1)}}) \\
 &\quad - \mathcal{L}_1(t, \mathcal{A}_{S_{(n-2)}})\| + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \left\| \int_0^t (\mathcal{L}_1(y, \mathcal{A}_{S_{(n-1)}}) \right. \\
 &\quad \left. - \mathcal{L}_1(y, \mathcal{A}_{S_{(n-2)}})) dy \right\|.
 \end{aligned}$$

Lipschitz condition leads us to

$$\begin{aligned}
 \|\mathcal{A}_{S_n}(t) - \mathcal{A}_{S_{(n-1)}}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 \|\mathcal{A}_{S_{(n-1)}} - \mathcal{A}_{S_{(n-2)}}\| \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 \times \int_0^t \|\mathcal{A}_{S_{(n-1)}} - \mathcal{A}_{S_{(n-2)}}\| dy.
 \end{aligned}$$

Next, we have

$$\begin{aligned}
 \|\hat{h}_{1n}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 \|\hat{h}_{1(n-1)}(t)\| \\
 &\quad + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 \int_0^t \|\hat{h}_{1(n-1)}(y)\| dy.
 \end{aligned} \tag{17}$$

Taking the same steps, we get

$$\begin{aligned}
 \|\hat{h}_{2n}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_2 \|\hat{h}_{2(n-1)}(t)\| + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_2 \int_0^t \|\hat{h}_{2(n-1)}(y)\| dy, \\
 \|\hat{h}_{3n}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_3 \|\hat{h}_{3(n-1)}(t)\| + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_3 \int_0^t \|\hat{h}_{3(n-1)}(y)\| dy, \\
 \|\hat{h}_{4n}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_4 \|\hat{h}_{4(n-1)}(t)\| + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_4 \int_0^t \|\hat{h}_{4(n-1)}(y)\| dy, \\
 \|\hat{h}_{5n}(t)\| &\leq 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_5 \|\hat{h}_{5(n-1)}(t)\| + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_5 \int_0^t \|\hat{h}_{5(n-1)}(y)\| dy,
 \end{aligned} \tag{18}$$

**Theorem 2** Exact coupled-solutions of the fractional-order model (3) of asthma disease exists if the below mentioned condition satisfies. That is one can find  $t_0$  in way that

$$2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 + 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 t_0 < 1.$$

**Proof.** As the state variables  $\mathcal{A}_S(t)$ ,  $\mathcal{A}_E(t)$ ,  $\mathcal{A}_I(t)$ ,  $\mathcal{A}_C(t)$ , and  $\mathcal{A}_P(t)$  are bounded. Moreover, we have shown that the Lipschitz condition is fulfilled by the kernels, Eqs. (17) and (18) gives the following by applying the recursive technique

$$\begin{aligned}
 \|\hat{h}_{1n}(t)\| &\leq \|\mathcal{A}_{S_n}(0)\| \left[ \left( 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 \right) + \left( 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 t \right) \right]^n, \\
 \|\hat{h}_{2n}(t)\| &\leq \|\mathcal{A}_{E_n}(0)\| \left[ \left( 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_2 \right) + \left( 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_2 t \right) \right]^n, \\
 \|\hat{h}_{3n}(t)\| &\leq \|\mathcal{A}_{I_n}(0)\| \left[ \left( 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_3 \right) + \left( 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_3 t \right) \right]^n, \\
 \|\hat{h}_{4n}(t)\| &\leq \|\mathcal{A}_{C_n}(0)\| \left[ \left( 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_4 \right) + \left( 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_4 t \right) \right]^n, \\
 \|\hat{h}_{5n}(t)\| &\leq \|\mathcal{A}_{P_n}(0)\| \left[ \left( 2 \frac{(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_5 \right) + \left( 2 \frac{\vartheta}{(2-\vartheta)U(\vartheta)} \mu_5 t \right) \right]^n.
 \end{aligned} \tag{19}$$

As a result, the existence of solutions of the asthma disease model and its continuity are achieved. In addition to this, we will show that the above is a solution of system (3), and proceed as follows

$$\begin{aligned}
 \mathcal{A}_S(t) - \mathcal{A}_S(0) &= \mathcal{A}_{S_n}(t) - \mathcal{M}1_n(t), \\
 \mathcal{A}_E(t) - \mathcal{A}_E(0) &= \mathcal{A}_{E_n}(t) - \mathcal{M}2_n(t), \\
 \mathcal{A}_I(t) - \mathcal{A}_I(0) &= \mathcal{A}_{I_n}(t) - \mathcal{M}3_n(t), \\
 \mathcal{A}_C(t) - \mathcal{A}_C(0) &= \mathcal{A}_{C_n}(t) - \mathcal{M}4_n(t), \\
 \mathcal{A}_P(t) - \mathcal{A}_P(0) &= \mathcal{A}_{P_n}(t) - \mathcal{M}5_n(t).
 \end{aligned}$$

Thus, we have

$$\begin{aligned}
 \|H_n(t)\| &= \left\| \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_1(t, \mathcal{A}_{S_n}) - \mathcal{L}_1(t, \mathcal{A}_{S_{(n-1)}})) \right. \\
 &\quad \left. + \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \times \int_0^t (\mathcal{L}_1(y, \mathcal{A}_{S_n}) - \mathcal{L}_1(y, \mathcal{A}_{S_{(n-1)}})) dy \right\|, \\
 &\leq \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \|\mathcal{L}_1(t, \mathcal{A}_{S_n}) - \mathcal{L}_1(t, \mathcal{A}_{S_{(n-1)}})\| \\
 &\quad + \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \times \int_0^t \|\mathcal{L}_1(y, \mathcal{A}_S) - \mathcal{L}_1(y, \mathcal{A}_{S_{(n-1)}})\| dy, \\
 &\leq \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 \|\mathcal{A}_{S_n} - \mathcal{A}_{S_{(n-1)}}\| + \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 \|\mathcal{A}_{S_n} - \mathcal{A}_{S_{(n-1)}}\| t.
 \end{aligned}$$

Following the technique, we proceed as

$$\| \mathcal{M}1_n(t) \| \leq \left( \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} + \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} t \right)^{n+1} \mu_1^{n+1} a.$$

Then the following is obtain at  $t_0$

$$\| \mathcal{M}1_n(t) \| \leq \left( \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} + \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} t_0 \right)^{n+1} \mu_1^{n+1} a. \quad (20)$$

The above Eq. (20) implies that

$$\| \mathcal{M}1_n(t) \| \rightarrow 0, \quad n \rightarrow \infty.$$

Following the same procedure, we obtain that  $\mathcal{M}2_n(t), \mathcal{M}3_n(t), \mathcal{M}4_n(t), \mathcal{M}5_n(t)$  approaches to 0 as  $n$  tends to  $\infty$ .

In the next step, we focus on the uniqueness of the solution of system (3), on contrast, we assume that  $(\mathcal{A}_{S1}(t), \mathcal{A}_{E1}(t), \mathcal{A}_{I1}(t), \mathcal{A}_{C1}(t), \mathcal{A}_{P1}(t))$  is another solution of system (3), then we have

$$\begin{aligned} \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) &= \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} (\mathcal{L}_1(t, \mathcal{A}_S) - \mathcal{L}_1(t, \mathcal{A}_{S1})) \\ &+ \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \times \int_0^t (\mathcal{L}_1(y, \mathcal{A}_S) - \mathcal{L}_1(y, \mathcal{A}_{S1})) dy. \end{aligned} \quad (21)$$


Applying the properties of norm, the above (21) converted into the following

$$\begin{aligned} \| \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) \| &\leq \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \| \mathcal{L}_1(t, \mathcal{A}_S) - \mathcal{L}_1(t, \mathcal{A}_{S1}) \| \\ &+ \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \times \int_0^t \| \mathcal{L}_1(y, \mathcal{A}_S) - \mathcal{L}_1(y, \mathcal{A}_{S1}) \| dy. \end{aligned}$$


Here, Lipschitz condition of kernel gives the following

$$\begin{aligned} \| \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) \| &\leq \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 \| \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) \| \\ &+ \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \times \int_0^t \mu_1 \| \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) \| dy, \end{aligned}$$


which gives




(a)



(b)



(c)



(d)

**Figure 2.** Illustration of the time series (solutions pathway) of susceptible, exposed, infected and smokers individuals of the proposed fractional-order model (3) of asthma disease with the variation of fractional-order  $\vartheta$ .

$$\| \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) \| \left( 1 - \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 - \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 t \right) \leq 0. \quad (22)$$

**Theorem 3** There exists a unique solution of the fractional-order model (3) of asthma disease if the below condition satisfied

$$\left( 1 - \frac{2(1-\vartheta)}{(2-\vartheta)U(\vartheta)} \mu_1 - \frac{2\vartheta}{(2-\vartheta)U(\vartheta)} \mu_1 t \right) > 0. \quad (23)$$

**Proof.** For the required result, we assume that the above condition (23) holds true then (22) implies that

$$\| \mathcal{A}_S(t) - \mathcal{A}_{S1}(t) \| = 0,$$

as a result of the uppermentioned, we have

$$\mathcal{A}_S(t) = \mathcal{A}_{S1}(t).$$


Following similar steps, we attain

$$\mathcal{A}_E(t) = \mathcal{A}_{E1}(t), \mathcal{A}_I(t) = \mathcal{A}_{I1}(t),$$

$$\mathcal{A}_C(t) = \mathcal{A}_{C1}(t), \mathcal{A}_P(t) = \mathcal{A}_{P1}(t).$$

### RESULTS AND DISCUSSION

We run multiple simulations in this portion of the study to better understand the complex dynamics of the suggested fractional-order asthma model. These simulations are necessary in order to find the most essential input factors that have a substantial impact on asthma prevalence in the general population. The fractional asthma system’s dynamical behaviour is quantitatively explored in order to present a more realistic picture of the illness. We assumed the following values of input parameters for simulation purposes:  $\beta = 0.0002$ ,  $\gamma = 0.00031$ ,  $\mu = 0.014$ ,  $\alpha = 0.258$ ,  $\lambda_1 = 0.0001$ ,  $\lambda = 0.00015$ ,  $\eta = 0.002$ ,  $\eta_1$




**Figure 3.** Illustration of dynamical behaviour of our proposed fractional-order model (3) of asthma disease with the variation of input parameter  $\eta_2$ .

= 0.0002,  $\eta_2 = 0.00025$  and  $q = 0.0001$  where the initial values of state-variables are given by:  $\mathcal{A}_S = 100$ ,  $\mathcal{A}_E = 50$ ,  $\mathcal{A}_I = 100$ ,  $\mathcal{A}_C = 10$  and  $\mathcal{A}_p = 50$  The time is taken in days in these numerical simulations.


Here, we perform four simulations to conceptualize the solution pathway of the susceptible, exposed, infected and smokers individuals of the asthma system and to show the influence of some input parameters on the dynamical behaviour of the system. In Figure 2, we illustrated the dynamics of asthma with the variation of fractional-order or index of memory  $\vartheta$ . We observed that the index of memory  $\vartheta$  has significance influence on solution pathway of asthma model and the control of  $\vartheta$  can highly control the dynamics of asthma disease. In Figure 3, we have shown the dynamical behaviour of the fractional system of asthma with the variation of the input parameter  $\eta_2$ , i.e.,  $\eta_2 = 0.00025, 0.00045, 0.00065, 0.00085$ , which shows that higher the value of  $\eta_2$  higher will be the level

of exposed and infected classes in the community. Thus, we can say that by controlling the smoker class can reduce the infection level of asthma in the community. In third simulation, we demonstrated the solution pathway of the proposed fractional-order model (3) of asthma with the variation of input parameter  $\lambda_1$ , i.e.,  $\lambda_1 = 0.0001, 0.0003, 0.0005, 0.0007$  shown in Figure 4. We noticed in second scenario that the increase of parameter  $\lambda_1$  highly decreases the level of  $\mathcal{A}_E$  while increase the level of asthma infected individuals  $\mathcal{A}_I$  in the community. In the last simulation presented in Figure 5, we again illustrated the solution pathway of the fractional asthma system with different values of fractional order, which implies that the endemic level of asthma can be decreased by lowering the fractional order  $\vartheta$ . Furthermore, we observed that the control of fractional-order can lower the level of infection, therefore, the index of memory  $\vartheta$  are suggested to the policy makers and medical experts to lower asthma attacks.



**Figure 4.** Illustration of the dynamical behaviour of the proposed fractional-order model (3) of asthma disease with the variation of input parameter  $\lambda_1$ .





**Figure 5.** Illustration of the time series (solutions pathway) of susceptible, exposed, infected and smokers individuals of the proposed fractional-order model (3) of asthma disease with the variation of fractional-order  $\vartheta$ .

**CONCLUSION**

Asthma is a well-known chronic disease that causes the inside walls of the airways of humans and make it hard to breathe. It is difficult to breath and talk in severe case of asthma which disturbs normal activity of humans. It is reported that smoking and environmental factors are critical and trigger asthma. In this article, we structured a mathematical model for asthma with environmental factors in the fractional derivative framework through CF fractional derivative. The suggested fractional model of asthma disease is then investigated for the basic properties through fractional calculus. The uniqueness and existence of the suggested asthma system are investigated via the theory of fixed point. Furthermore, we represent a novel numerical technique to highlight the time series of fractional-order asthma model. We have shown the dynamical behaviour of different stages of asthma model with variation of

fractional-order  $\vartheta$ . It is predicted that the index of memory  $\vartheta$  has a positive effect on the system and can be used as a control parameter. Finally, the dynamical behaviour of asthma with input factors are then highlighted numerically to show the influence of several input parameter on the time series of asthma. We suggested to the policy makers that the index of memory or fractional-order can play an important role in the control of asthma. We will further study the epidemiology of asthma to formulate a more comprehensive model for the transmission phenomena of asthma to produce more accurate and precious results with real data in the future work. Moreover, sensitivity test through PRCC method will be performed to determine the most critical input factor for the control and prevention of asthma. We will also introduce some control policies through optimal control theory and impulsive vaccination for the prevention of asthma infection in the society.

## AUTHORSHIP CONTRIBUTIONS

Authors equally contributed to this work.

## DATA AVAILABILITY STATEMENT

The authors confirm that the data that supports the findings of this study are available within the article. Raw data that support the finding of this study are available from the corresponding author, upon reasonable request.

## CONFLICT OF INTEREST

The author declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## ETHICS

There are no ethical issues with the publication of this manuscript.

## REFERENCES

- [1] Ghosh M. Industrial pollution and Asthma: A mathematical model. *J Biol Syst* 2000;8:347–371. [\[CrossRef\]](#)
- [2] Polosa R, Thomson NC. Smoking and asthma: dangerous liaisons. *Eur Respir J* 2013;41:716–726. [\[CrossRef\]](#)
- [3] Strachan DP. The role of environmental factors in asthma. *Br Med Bull* 2000;56:865–882. [\[CrossRef\]](#)
- [4] Driscoll AJ, Arshad SH, Bont L, Brunwasser SM, Cherian T, Englund JA, et al. Does respiratory syncytial virus lower respiratory illness in early life cause recurrent wheeze of early childhood and asthma? Critical review of the evidence and guidance for future studies from a World Health Organization-sponsored meeting. *Vaccine* 2020;38:2435–2448. [\[CrossRef\]](#)
- [5] Canadian Medical Association. Secondhand cigarette smoke worsens symptoms in children with asthma. Section on Allergy, Canadian Paediatric Society. *CMAJ* 1986;135:321–323.
- [6] Collishaw NE, Kirkbride J, Wigle DT. Tobacco smoke in the workplace: an occupational health hazard. *Can Med Assoc J* 1984;131:1199–1204.
- [7] Dockery DW, Pope CA, Xu X, Spengler JD, Ware JH, Fay ME, et al. An association between air pollution and mortality in six US cities. *N Eng J Med* 1993;329:1753–1759. [\[CrossRef\]](#)
- [8] Utell MJ, Looney RJ. Environmentally induced asthma. *Toxicol Lett* 1995;82:47–53. [\[CrossRef\]](#)
- [9] Jan R, Xiao Y. Effect of partial immunity on transmission dynamics of dengue disease with optimal control. *Math Methods Appl Sci* 2019;42:1967–1983. [\[CrossRef\]](#)
- [10] Secer A, Onder I, Ozışık M. Sinc-Galerkin method for solving system of singular perturbed reaction-diffusion problems. *Sigma J Eng Nat Sci* 2021;39:203–212. [\[CrossRef\]](#)
- [11] Sisman S, Merdan M. Global stability of Susceptible Diabetes Complication (SDC) model in discrete time. *Sigma J Eng Nat Sci* 2021;39:290–312. [\[CrossRef\]](#)
- [12] Ergene B, Yalçın B. A finite element study on modal analysis of lightweight pipes. *Sigma J Eng Nat Sci* 2021;39:268–278. [\[CrossRef\]](#)
- [13] Akbari N, Gholinia M, Gholinia S, Dabbaghian S, Ganji DD. Analytical and numerical study of hydrodynamic nano fluid flow in a tow-dimensional semi-porous channel with transverse magnetic field. *Sigma J Eng Nat Sci* 2018;36:587–608.
- [14] Jan R, Xiao Y. Effect of pulse vaccination on dynamics of dengue with periodic transmission functions. *Adv Differ Equ* 2019;2019:1–7. [\[CrossRef\]](#)
- [15] Jan R, Khan MA, Gómez-Aguilar JF. Asymptomatic carriers in transmission dynamics of dengue with control interventions. *Optim Control Appl Methods* 2020;41:430–447. [\[CrossRef\]](#)
- [16] Attaullah, Jan R, and Yüzbaşı S. Dynamical behaviour of HIV Infection with the influence of variable source term through Galerkin method. *Chaos Solit Fractals* 2021;152:111429. [\[CrossRef\]](#)
- [17] Mahmoudi MR, Baleanu D, Band SS, Mosavi A. Factor analysis approach to classify COVID-19 datasets in several regions. *Results Phys* 2021;25:104071. [\[CrossRef\]](#)
- [18] Yüzbaşı S. A numerical approach to solve the model for HIV infection of CD4+ T cells. *Appl Math Model* 2012;36:976352. [\[CrossRef\]](#)
- [19] Yüzbaşı S, Ismailov N. A numerical method for the solutions of the HIV infection model of CD4+ T-cells. *Int J Biomath* 2017;10:1750098. [\[CrossRef\]](#)
- [20] Ahmed N, Rafiq M, Adel W, Rezazadeh H, Khan I, Nisar KS. Structure preserving numerical analysis of HIV and CD4+ T-cells reaction diffusion model in two space dimensions. *Chaos Solit Fractals* 2020;139:110307. [\[CrossRef\]](#)
- [21] Thirumalai S, Seshadri R, Yüzbaşı S. On the solution of the Human Immunodeficiency Virus (HIV) infection model using spectral collocation method. *Int J Biomath* 2021;14:2050074. [\[CrossRef\]](#)
- [22] Yüzbaşı S. An exponential collocation method for the solutions of the HIV infection model of CD4+ T cells. *Int J Biomath* 2016;9:1650036. [\[CrossRef\]](#)
- [23] Winkler T, Venegas JG, Harris RS. Mathematical modeling of ventilation defects in asthma. *Drug Discov Today Dis Models* 2015;15:3–8. [\[CrossRef\]](#)
- [24] Joseph GA, Balamuralitharan S. A Nonlinear differential equation model of Asthma effect of environmental pollution using LHAM. In *Journal of Physics: Conference Series* 2018; (Vol. 1000, No. 1, p. 012043). IOP Publishing. [\[CrossRef\]](#)

- [25] Rosenberg HF, Druey KM. Modeling asthma: Pitfalls, promises, and the road ahead. *J Leukoc Biol* 2018;104:41–48. [\[CrossRef\]](#)
- [26] Burrowes KS, Doel T, Brightling C. Computational modeling of the obstructive lung diseases asthma and COPD. *J Transl Med* 2014;12:1–8. [\[CrossRef\]](#)
- [27] Sheffield PE, Knowlton K, Carr JL, Kinney PL. Modeling of regional climate change effects on ground-level ozone and childhood asthma. *Am J Prev Med* 2011;41:251–257. [\[CrossRef\]](#)
- [28] Jang Y, Shin H, Lee MK, Kwon OS, Shin JS, Kim YI, Kim CW, Lee HR, Kim M. Antiviral activity of lambda-carrageenan against influenza viruses and severe acute respiratory syndrome coronavirus 2. *Sci Rep* 2021;11:1–2. [\[CrossRef\]](#)
- [29] Slowikowska M, Bajzert J, Miller J, Stefaniak T, Niedzwiedz A. The dynamics of circulating immune complexes in horses with severe equine asthma. *Animals* 2021;11:1001. [\[CrossRef\]](#)
- [30] Nápoles JEN, Guzmán PM, Lugo LM, Kashuri, A. The local generalized derivative and Mittag Leffler function. *Sigma J Eng Nat Sci* 2020;38:1007–1017.
- [31] Shah Z, Jan R, Kumam P, Deebani W, Shutaywi M. Fractional dynamics of HIV with source term for the supply of new CD4+ T-cells depending on the viral load via caputo-fabrizio derivative. *Molecules* 2021;26:1806. [\[CrossRef\]](#)
- [32] Srivastava HM, Jan R, Jan A, Deebani W, Shutaywi M. Fractional-calculus analysis of the transmission dynamics of the dengue infection. *Chaos* 2021;31:053130. [\[CrossRef\]](#)
- [33] Jan R, Jan A. MSGDTM for solution of fractional order dengue disease model. *Int J Sci Res* 2017;6:1140–1144.
- [34] Fatmawati F, Jan R, Khan MA, Khan Y, Ullah S. A new model of dengue fever in terms of fractional derivative. *Math Biosci Eng* 2020;17:5267–5287. [\[CrossRef\]](#)
- [35] Podlubny I. Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. 1st ed. Amsterdam: Elsevier; 1998.
- [36] Samko SG, Kilbas AA, Marichev OI. Fractional integrals and derivatives, translated from the 1987 Russian original. Switzerland ; Philadelphia, Pa., USA : Gordon and Breach Science Publishers; 1993.
- [37] Qureshi S. Periodic dynamics of rubella epidemic under standard and fractional Caputo operator with real data from Pakistan. *Math Comput Simul* 2020;178:151–165. [\[CrossRef\]](#)
- [38] Qureshi S. Effects of vaccination on measles dynamics under fractional conformable derivative with Liouville-Caputo operator. *Eur Phys J Plus* 2020;135:1–20. [\[CrossRef\]](#)
- [39] Qureshi S, Rangaig NA, Baleanu D. New numerical aspects of Caputo-Fabrizio fractional derivative operator. *Mathematics* 2019;7:374. [\[CrossRef\]](#)
- [40] Jan R, Khan MA, Kumam P, Thounthong P. Modeling the transmission of dengue infection through fractional derivatives. *Chaos Solit Fractals* 2019;127:189–216. [\[CrossRef\]](#)
- [41] Qureshi S, Yusuf A. Fractional derivatives applied to MSEIR problems: Comparative study with real world data. *Eur Phys J Plus* 2019;134:1–3. [\[CrossRef\]](#)
- [42] Jan A, Jan R, Khan H, Zobaer MS, Shah R. Fractional-order dynamics of Rift Valley fever in ruminant host with vaccination. *Commun Math Biol Neurosci* 2020;2020:79.
- [43] Jan R, Khan H, Kumam P, Tchier F, Shah R, Bin Jebreen H. The investigation of the fractional-view dynamics of helmholtz equations within caputo operator. *Comput Mater Contin* 2021;68:3185–3201. [\[CrossRef\]](#)
- [44] Muresan CI, Birs IR, Dulf EH. Event-based implementation of fractional order IMC controllers for simple FOPDT processes. *Mathematics* 2020;8:1378. [\[CrossRef\]](#)
- [45] Goufo EF. A biomathematical view on the fractional dynamics of cellulose degradation. *Fract Calc Appl Anal* 2015;18:554–564. [\[CrossRef\]](#)
- [46] Yusuf A, Qureshi S, Shah SF. Mathematical analysis for an autonomous financial dynamical system via classical and modern fractional operators. *Chaos Solit Fractals* 2020;132:109552. [\[CrossRef\]](#)
- [47] Sweilam NH, Al-Mekhlafi SM, Baleanu D. A hybrid stochastic fractional order Coronavirus (2019-nCov) mathematical model. *Chaos Solit Fractals* 2021;145:110762. [\[CrossRef\]](#)
- [48] Caputo M, Fabrizio M. A new definition of fractional derivative without singular kernel. *Progr Fract Differ Appl* 2015;1:1–3. [\[CrossRef\]](#)
- [49] Losada J, Nieto JJ. Properties of a new fractional derivative without singular kernel. *Progr Fract Differ Appl* 2015;1:87–92.
- [50] Attaullah, Sohaib M. Mathematical modeling and numerical simulation of HIV infection model. *Results Appl Math* 2020;7:100118. [\[CrossRef\]](#)
- [51] Attaullah, Jan R, Jabeen A. Solution of the HIV infection model with full logistic proliferation and variable source term using Galerkin scheme. *Matrix Sci Math* 2020;4:37–43.
- [52] Attaullah, Zeeshan, Tufail Khan M, Alyobi S, Yassen MF, Prathumwan D. A computational approach to a model for HIV and the immune system interaction. *Axioms* 2022;11:578. [\[CrossRef\]](#)
- [53] Attaullah RD, Khurshaid A, Alyobi S, Yassen MF, Prathumwan D. Computational framework of the SVIR epidemic model with a non-linear saturation incidence rate. *Axioms* 2022;11:651. [\[CrossRef\]](#)

- [54] Attaullah RD, Weera W. Galerkin time discretization scheme for the transmission dynamics of HIV infection with non-linear supply rate. *J AIMS Math* 2022;6:11292–11310. [[CrossRef](#)]
- [55] Attaullah RD, Alyobi S, Yassen MF. A study on the transmission and dynamical behavior of an HIV/AIDS epidemic model with a cure rate. *AIMS Math* 2022;7:17507–17528. [[CrossRef](#)]
- [56] Attaullah RD, Yassen MF, Alyobi S, Al-Duais FS, Weera W. On the comparative performance of fourth order Runge-Kutta and the Galerkin-Petrov time discretization methods for solving nonlinear ordinary differential equations with application to some mathematical models in epidemiology. *AIMS Math* 2023;8:3699–3729. [[CrossRef](#)]
- [57] Attaullah, Jawad M, Alyobi S, Yassen MF, Weera W. A higher order Galerkin time discretization scheme for the novel mathematical model of COVID-19. *AIMS Math* 2023;8:3763–3790. [[CrossRef](#)]
- [58] Attaullah, Yüzbaşı Ş, Alyobi S, Yassen MF, Weera W. A Higher-Order Galerkin Time Discretization and Numerical Comparisons for Two Models of HIV Infection. *Comput Math Methods Med* 2022;2022:3599827. [[CrossRef](#)]