



Research Article

## A consensus reaching process with fuzzy matrix of interpersonal influences

Asma MAHMOOD<sup>1,\*</sup>

<sup>1</sup>Department of Mathematics, GC University, Faisalabad, 38000, Pakistan

### ARTICLE INFO

#### Article history

Received: 04 December 2023

Revised: 15 February 2024

Accepted: 04 May 2024

#### Keywords:

Fuzzy Inverse Matrix; Group Decision Making; Influence Model; Triangular Fuzzy Numbers

### ABSTRACT

Consensus reaching process is an influential feature of social influence network theory and group decision making. The influence model undertakes a pivotal role where group decision making is based on opinion dynamics. Interpersonal influences of experts have an exceptional part of the influence model which is a dominant addition in social influence networks. Influence based model is a simple and satisfactory mathematical representation of the change of opinions due to the experts' influences. Uncertainty is presupposed in almost every direction of decision making and opinion dynamics so should also be included in the influence model. In this paper, the influence model is refined by utilizing the triangular fuzzy numbers in place of crisp numbers, where not only the initial opinions but also the interpersonal influences are represented as fuzzy numbers. This extends the influence model from ordinary numbers to fuzzy numbers. A fuzzy inverse matrix is computed by using a system of linear equations where coefficients and constants are fuzzy numbers. These equations are called fuzzy linear equations.

**Cite this article as:** Mahmood A. A consensus reaching process with fuzzy matrix of interpersonal influences. Sigma J Eng Nat Sci 2025;43(2):607–614.

### INTRODUCTION

Decision-making is an imperative element of our routine life whether we are at home or work and have become complicated due to the development of society. Organizations arrange group of members for their decision-making processes, which is known as group decision making (GDM). GDM models are usually concerned with two processes: the consensus reaching process (CRP) [1-3] and the selection process [4,5]. CRP is an iterative process leading to the final solution after discussions and interpersonal influences of experts. Yao and Gu [6] propose a consensus model based on

an influence network for large-scale GDM. After the influence model was introduced by Friedkin and Johnsen [7], there have been presented its related concepts and applications by different authors [8-10]. Influence models contribute particularly in GDM, where initial opinions of  $n$  experts are revised due to their interpersonal and social influences [5,11]. These models are based on an iterative process where the matrix  $W = [w_{ij}]_{n \times n}$  ( $w_{ij} \in [0,1]$ ), of interpersonal influences is a basic component. Weights  $w_{ij}$  are assumed to be satisfy the normalization property i.e.  $\sum_{j=1}^n w_{ij} = 1$ .

Fuzzy set theory [12] is suggested to deal with uncertainty which is an important aspect of almost all

#### \*Corresponding author.

\*E-mail address: [asmamahmood@gcuf.edu.pk](mailto:asmamahmood@gcuf.edu.pk)

This paper was recommended for publication in revised form by Editor-in-Chief Ahmet Selim Dalkilic



decision-making processes. Fuzzy set have many generalizations with a large number of applications in optimization and other decision making [13-15]. The mathematical theory of interpersonal influences leads us to construct the influence models. Fuzzy preference relations (FPRs) also play an important role in such problems. FPRs are utilized to form the initial opinions of experts and through a mathematical model final opinions are obtained [16-18].

Fuzzy numbers have been introduced by Jain [19] and Dubois and Prade [20] to model the uncertain information. Ranking fuzzy numbers is a crucial step in investigating fuzzy information in decision making process. A lot of contributions have been made in ranking of fuzzy numbers [21-24]. A fuzzy number gets its extension from the real number and can be considered as a function whose domain is a specified set. A fuzzy set satisfying some conditions, is called a fuzzy number. Fuzzy numbers and the extension principle are the basis for fuzzy arithmetic [25]. Lee [26] has also discussed fuzzy numbers with their basic operations  $\oplus$ ,  $\ominus$ ,  $\otimes$  and  $\oslash$ , where these operations are conducted for triangular fuzzy numbers (TFNs), with the help of  $\alpha$ -cuts. Such operations are not fuzzy, the numbers on which the operations are performed are fuzzy, so the results of these operations are also fuzzy. TFN  $(a_l, a_0, a_r)$  assigns an interval, called  $\alpha$ -cut, for each value in the interval  $[0,1]$ . Fuzzy numbers have a wide utilization in decision sciences and engineering applications [27,28].

A fuzzy matrix has two different meanings in the literature: firstly,  $A = (a_{ij})_{m \times n}$  is called a fuzzy matrix if  $a_{ij} \in [0,1]$ , ( $i = 1, 2, \dots, m; j = 1, 2, \dots, n$ ), on the other hand a matrix with entries of fuzzy numbers is called a fuzzy matrix too. Fuzzy matrices of second class is considered in this article. There is a strong connection between interval matrix and fuzzy matrix because each  $\alpha$ -cut of a fuzzy matrix is an interval matrix that contains interval numbers. The interval matrix, its operations and singularity were discussed by Rohn [29,30]. Clustering with covariance matrix and mediative fuzzy relations which contain a large number of uncertainties, also took advantage of fuzzy matrices [31-34]. An approximate inverse of an uncertain matrix was introduced in Ghaoui [35]. A square matrix  $A$  of order  $n$  represents the co-efficient matrix in a system  $Ax = b$  of linear equations whose solution is uniquely determined by finding the inverse of  $A$ . In the case of a fuzzy matrix, the computation of its inverse is quite complicated. Dehghan et al. [36] presented some conditions for the invertibility of fuzzy matrices in terms of interval matrices. Farahani et.al. [37] presented a method to find the inverse of a fuzzy matrix by using eigen value method. Dequan and Guo [38] investigated a class of fuzzy linear matrix equation by using the embedding approach. Some recent developments are observed in neutrosophic fuzzy matrices [39,40]. Basaran [41] has suggested a method to calculate fuzzy inverse matrix by using a fuzzy linear equation

system. This method of obtaining the inverse fuzzy matrix is utilized in this paper for the matrix of triangular fuzzy numbers and to present the influence model where the matrix of interpersonal influences is a fuzzy matrix with triangular fuzzy numbers as its entries.

The concepts of social influence network theory and opinion dynamics are not mathematical in nature, but formation of the matrices of opinions and influences prepares the fundamentals for mathematical procedures. This research aims to distend the insufficiency of mathematical techniques in opinion dynamics and social influences. Crisp numbers are replaced by fuzzy numbers in the CRP [7]. TFNs are utilized which are suitable for ranking and comparison of the influences of experts. TFNs give comprehensive results while inverse of the matrix is computed.

## PRELIMINARIES

### Triangular Fuzzy Numbers

#### Definition 1

[27] A fuzzy number  $A = (a_l, a_0, a_r)$  where  $a_l, a_0, a_r \in R$  and  $a_l \leq a_0 \leq a_r$ , is a triangular fuzzy number if it is represented as the following membership function:

$$\mu_{(A)}(x) = \begin{cases} 0, & x < a_l \\ \frac{x - a_l}{a_0 - a_l} & a_l \leq x \leq a_0 \\ \frac{a_r - x}{a_r - a_0} & a_0 \leq x \leq a_r \\ 0 & x > a_r \end{cases}$$

### Operations on fuzzy numbers

The arithmetic operations on fuzzy numbers can be defined by the extension principle.

#### Definition 2

[43] Let  $A$  and  $B$  be two fuzzy numbers and  $\times$  be an operation on  $R$ , such as  $+$ ,  $-$ ,  $*$ ,  $\div$ ,  $\dots$ . By extension principle, the extended operation  $\otimes$  on fuzzy numbers can be defined by:

$$\mu_{A \otimes B}(z) = \sup_{x, y: z = x \times y} \min(A(x), B(y))$$

Suppose  $A = (a_l, a_0, a_r)$  and  $B = (b_l, b_0, b_r)$  are two triangular fuzzy numbers. The formulae for the extended addition, subtraction, multiplication and division operations become [17,23]:

#### 1. Addition

$$\begin{aligned} A \oplus B &= (a_l, a_0, a_r) \oplus (b_l, b_0, b_r) \\ &= (a_l + b_l, a_0 + b_0, a_r + b_r). \end{aligned}$$

#### 2. Subtraction

$$\begin{aligned} A \ominus B &= (a_l, a_0, a_r) \ominus (b_l, b_0, b_r) \\ &= (a_l - b_l, a_0 - b_0, a_r - b_r). \end{aligned}$$

**3. Multiplication**

$$A \otimes B \cong (a_l, a_0, a_r) \otimes (b_l, b, b_r) \cong (a_l b_l, a_0 b_0, a_r b_r).$$

**4. Division**

$$A \oslash B \cong (a_l, a_0, a_r) \oslash (b_l, b, b_r) \cong \left( \frac{a_l}{b_r}, \frac{a_0}{b_0}, \frac{a_r}{b_l} \right).$$

**Ordering of triangular fuzzy numbers**

[43] The set of all fuzzy numbers is denoted by  $N_F$ . A total ordering  $\leq$  on  $N_T$  (the set of all triangular fuzzy numbers) may be defined as:  $(a_l, a_0, a_r) \leq (b_l, b, b_r)$  if and only if

1.  $a_l < b_l$ , or
2.  $a_l = b_l$  but  $a_0 < b_0$ , or
3.  $a_l = b_l, a_0 = b_0$  but  $a_r < b_r$ .

**Fuzzy Inverse Matrix**

**Definition 3**

[44] A matrix  $\tilde{A} = (\tilde{a}_{ij})$  is called a fuzzy matrix, if each element of  $\tilde{A}$  is a fuzzy number.

let  $A = (a_l, a_0, a_r)$  and  $B = (b_l, b_0, b_r)$  be two  $m \times n$  and  $n \times p$  fuzzy matrices. The size of the product of two fuzzy matrices is  $m \times p$  and is written as follows:  $\tilde{A} \times \tilde{B} = \tilde{C} = (\tilde{c}_{ij})$ , where  $\tilde{c}_{ij} = \oplus(\tilde{a}_{ik} \otimes \tilde{b}_{kj})$  where  $\otimes$  is the approximated multiplication.

Before defining fuzzy inverse matrix, it is necessary to define fuzzy zero number and fuzzy one number to develop fuzzy identity matrix.

**Definition 4**

[41] If the center value of a fuzzy number is 0 and the left and right spread values are  $\alpha$  and  $\beta$  where  $0 < \alpha < \beta < 1$ , this fuzzy number is called fuzzy zero number and is denoted as  $\tilde{0} = (-\alpha, 0, \beta)$ .

**Definition 5**

If the center value of a fuzzy number is  $\tilde{1}$  and the left and right spread values are  $\delta$  and  $\lambda$  where  $0 < \delta < \lambda < 1$ , this

fuzzy number is called fuzzy one number and is denoted as  $\tilde{1} = (1 - \delta, 1, 1 + \lambda)$ .

**Definition 6**

If the diagonal elements of a fuzzy matrix are fuzzy one numbers and the off-diagonal elements are fuzzy zero numbers, then this fuzzy matrix is called fuzzy identity matrix and is denoted by  $\tilde{I}$ .

$$\tilde{I} = \begin{bmatrix} \tilde{1} & \tilde{0} & \dots & \tilde{0} \\ \tilde{0} & \tilde{1} & \dots & \tilde{0} \\ \vdots & \vdots & \ddots & \vdots \\ \tilde{0} & \tilde{0} & \dots & \tilde{1} \end{bmatrix}.$$

**Social Influence Network Theory**

Let  $W = w_{ij}$  be the matrix of interpersonal influences among  $n$  experts with  $w_{ij} \in [0,1]$  and  $\sum_{j=1}^n w_{ij} = 1$ . The diagonal matrix  $A = \text{diag}(a_{11}, a_{22}, \dots, a_{nn})$  is obtained from the matrix  $W$  computing  $a_{ii} = 1 - w_{ii}, i = 1, 2, \dots, n$ . The matrix  $A$  is clarified as the susceptibility of all experts to interpersonal influence [8]. With the first opinion  $g^{(1)}$ , the following iterative plan is suggested to find the revised and final opinion:

$$g^{(t)} = AWg^{(t-1)} + (I - A)g^{(1)} \tag{1}$$


If  $I - AW$  is non-singular, then this process reaches the following

$$g^{(\infty)} = (I - AW)^{-1}(I - A)g^{(1)}$$

Figure 1 describes the general consensus reaching process with interpersonal influences of experts.

**INFLUENCE MODEL WITH FUZZY MATRICES**

Fuzzy numbers generalize real numbers and are very useful to represent data corresponding to uncertain situations. In this section, an influence model is presented which is similar to the model presented in section 2.3 but the ordinary numbers are replaced by fuzzy numbers. TFNs consider only 3 data points and two linear functions so suitable to represent the influences and opinions. Final opinions are



**Figure 1.** Consensus reaching process.

obtained when initial opinions and IIs of experts are given. Stepwise procedure is as follows:

**Step 1:**

A fuzzy matrix

$$\tilde{W} = \begin{bmatrix} \tilde{w}_{11} & \tilde{w}_{12} & \dots & \tilde{w}_{1n} \\ \tilde{w}_{21} & \tilde{w}_{22} & \dots & \tilde{w}_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \tilde{w}_{n1} & \tilde{w}_{n2} & \dots & \tilde{w}_{nn} \end{bmatrix}$$

is taken as a matrix of interpersonal influences, where  $\tilde{w}_{ij}$  are assumed to be triangular fuzzy numbers and

$\tilde{w}_{i1} \oplus \tilde{w}_{i2} \oplus \dots \oplus \tilde{w}_{in} = \tilde{1}$  for  $i = 1, 2, \dots, n$ . And

$$\tilde{A} = \begin{bmatrix} \tilde{a}_{11} & 0 & \dots & 0 \\ 0 & \tilde{a}_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \tilde{a}_{nn} \end{bmatrix}$$

is a matrix of sensitivities of experts to the interpersonal influences,  $\tilde{a}_{ii} = \tilde{1} - \tilde{w}_{ii}$ .

**Step 2:**

A fuzzy column matrix

$$\tilde{y}^{(1)} = \begin{pmatrix} \tilde{y}_1^{(1)} \\ \tilde{y}_2^{(1)} \\ \vdots \\ \tilde{y}_n^{(1)} \end{pmatrix}$$

is taken where its entries are the initial opinions of n experts.

**Step 3:**

Compute  $\tilde{A}\tilde{W}$

$$\tilde{A}\tilde{W} = \begin{bmatrix} \tilde{w}_{11} \otimes \tilde{a}_{11} & \tilde{w}_{12} \otimes \tilde{a}_{22} & \dots & \tilde{w}_{1n} \otimes \tilde{a}_{nn} \\ \tilde{w}_{21} \otimes \tilde{a}_{11} & \tilde{w}_{22} \otimes \tilde{a}_{22} & \dots & \tilde{w}_{2n} \otimes \tilde{a}_{nn} \\ \vdots & \vdots & \ddots & \vdots \\ \tilde{w}_{n1} \otimes \tilde{a}_{11} & \tilde{w}_{n2} \otimes \tilde{a}_{22} & \dots & \tilde{w}_{nn} \otimes \tilde{a}_{nn} \end{bmatrix},$$

then  $\tilde{I} - \tilde{A}\tilde{W}$  and  $\tilde{I} - \tilde{A}$  by using  $\tilde{I}$  as defined in Definition 4.

**Step 4:**

Find  $(\tilde{I} - \tilde{A}\tilde{W})^{-1}$  by the procedure described in section 3.1 and then final opinions  $y^\infty$  by using the following equation:

$$y^{(\infty)} = (\tilde{I} - \tilde{A}\tilde{W})^{-1}(\tilde{I} - \tilde{A})y^{(1)}$$

**Fuzzy Inverse Matrix**

Following is a step-wise procedure to find fuzzy inverse of a fuzzy matrix with the entries represented as triangular fuzzy numbers:

**Step 1:**

Let  $\tilde{X} = \begin{bmatrix} \tilde{x}_{11} & \tilde{x}_{12} & \tilde{x}_{13} \\ \tilde{x}_{21} & \tilde{x}_{22} & \tilde{x}_{23} \\ \tilde{x}_{31} & \tilde{x}_{32} & \tilde{x}_{33} \end{bmatrix}$

be a matrix with entries of triangular fuzzy numbers  $\tilde{x}_{ij} = (x_{ij} - u_{ij}^L, x_{ij}, x_{ij} + u_{ij}^R)$ ,  $i, j = 1, 2, 3$ .  $u_{ij}^L$  and  $u_{ij}^R$  represent left and right spreads of  $\tilde{x}_{ij}$  respectively. Let

$$\tilde{Z} = \begin{bmatrix} \tilde{z}_{11} & \tilde{z}_{12} & \tilde{z}_{13} \\ \tilde{z}_{21} & \tilde{z}_{22} & \tilde{z}_{23} \\ \tilde{z}_{31} & \tilde{z}_{32} & \tilde{z}_{33} \end{bmatrix}$$

be the inverse of  $\tilde{X}$  where  $z_{ij} = (z_{ij} - v_{ij}^L, z_{ij}, z_{ij} + v_{ij}^R)$ ,  $i, j = 1, 2, 3$ .  $v_{ij}^L$  and  $v_{ij}^R$  represent left and right spreads of  $\tilde{z}_{ij}$  respectively. Then  $\tilde{X} \otimes \tilde{Z} = \tilde{I}$ .

**Step 2:**

We will solve the following system of equations:

$$\begin{aligned} (\tilde{x}_{11} \otimes \tilde{z}_{11}) \oplus (\tilde{x}_{12} \otimes \tilde{z}_{21}) \oplus (\tilde{x}_{13} \otimes \tilde{z}_{31}) &= \tilde{1}, \\ (\tilde{x}_{11} \otimes \tilde{z}_{12}) \oplus (\tilde{x}_{12} \otimes \tilde{z}_{22}) \oplus (\tilde{x}_{13} \otimes \tilde{z}_{32}) &= \tilde{0}, \\ (\tilde{x}_{11} \otimes \tilde{z}_{13}) \oplus (\tilde{x}_{12} \otimes \tilde{z}_{23}) \oplus (\tilde{x}_{13} \otimes \tilde{z}_{33}) &= \tilde{0}, \\ (\tilde{x}_{21} \otimes \tilde{z}_{11}) \oplus (\tilde{x}_{22} \otimes \tilde{z}_{21}) \oplus (\tilde{x}_{23} \otimes \tilde{z}_{31}) &= \tilde{0}, \\ (\tilde{x}_{21} \otimes \tilde{z}_{12}) \oplus (\tilde{x}_{22} \otimes \tilde{z}_{22}) \oplus (\tilde{x}_{23} \otimes \tilde{z}_{32}) &= \tilde{1}, \\ (\tilde{x}_{21} \otimes \tilde{z}_{13}) \oplus (\tilde{x}_{22} \otimes \tilde{z}_{23}) \oplus (\tilde{x}_{23} \otimes \tilde{z}_{33}) &= \tilde{0}, \\ (\tilde{x}_{31} \otimes \tilde{z}_{11}) \oplus (\tilde{x}_{32} \otimes \tilde{z}_{21}) \oplus (\tilde{x}_{33} \otimes \tilde{z}_{31}) &= \tilde{0}, \\ (\tilde{x}_{31} \otimes \tilde{z}_{12}) \oplus (\tilde{x}_{32} \otimes \tilde{z}_{22}) \oplus (\tilde{x}_{33} \otimes \tilde{z}_{32}) &= \tilde{0}, \\ (\tilde{x}_{31} \otimes \tilde{z}_{13}) \oplus (\tilde{x}_{32} \otimes \tilde{z}_{23}) \oplus (\tilde{x}_{33} \otimes \tilde{z}_{33}) &= \tilde{1}. \end{aligned}$$

**Step 3:**

Following system is solved for center part  $z_{ij}$  of the inverse matrix:

$$\begin{aligned} x_{11}z_{11} + x_{12}z_{21} + x_{13}z_{31} &= 1, \\ x_{11}z_{12} + x_{12}z_{22} + x_{13}z_{32} &= 0, \\ x_{11}z_{13} + x_{12}z_{23} + x_{13}z_{33} &= 0, \\ x_{21}z_{11} + x_{22}z_{21} + x_{23}z_{31} &= 0, \\ x_{21}z_{12} + x_{22}z_{22} + x_{23}z_{32} &= 1, \\ x_{21}z_{13} + x_{22}z_{23} + x_{23}z_{33} &= 0, \\ x_{31}z_{11} + x_{32}z_{21} + x_{33}z_{31} &= 0, \\ x_{31}z_{12} + x_{32}z_{22} + x_{33}z_{32} &= 0, \\ x_{31}z_{13} + x_{32}z_{23} + x_{33}z_{33} &= 1. \end{aligned}$$

To find the left spread part  $v_{ij}^L$  of the matrix, system is written as follows:

$$\begin{aligned} x_{11}v_{11}^L + u_{11}^Lz_{11} + x_{12}v_{21}^L + u_{12}^Lz_{21} + x_{13}v_{31}^L + u_{13}^Lz_{31} &= \delta, \\ x_{11}v_{12}^L + u_{11}^Lz_{12} + x_{12}v_{22}^L + u_{12}^Lz_{22} + x_{13}v_{32}^L + u_{13}^Lz_{32} &= \alpha, \\ x_{11}v_{13}^L + u_{11}^Lz_{13} + x_{12}v_{23}^L + u_{12}^Lz_{23} + x_{13}v_{33}^L + u_{13}^Lz_{33} &= \alpha, \\ x_{21}v_{11}^L + u_{21}^Lz_{11} + x_{22}v_{21}^L + u_{22}^Lz_{21} + x_{23}v_{31}^L + u_{23}^Lz_{31} &= \alpha, \\ x_{21}v_{12}^L + u_{21}^Lz_{12} + x_{22}v_{22}^L + u_{22}^Lz_{22} + x_{23}v_{32}^L + u_{23}^Lz_{32} &= \delta, \\ x_{21}v_{13}^L + u_{21}^Lz_{13} + x_{22}v_{23}^L + u_{22}^Lz_{23} + x_{23}v_{33}^L + u_{23}^Lz_{33} &= \alpha, \\ x_{31}v_{11}^L + u_{31}^Lz_{11} + x_{32}v_{21}^L + u_{32}^Lz_{21} + x_{33}v_{31}^L + u_{33}^Lz_{31} &= \alpha, \\ x_{31}v_{12}^L + u_{31}^Lz_{12} + x_{32}v_{22}^L + u_{32}^Lz_{22} + x_{33}v_{32}^L + u_{33}^Lz_{32} &= \alpha, \\ x_{31}v_{13}^L + u_{31}^Lz_{13} + x_{32}v_{23}^L + u_{32}^Lz_{23} + x_{33}v_{33}^L + u_{33}^Lz_{33} &= \delta. \end{aligned}$$

Similarly, right spread part  $v_{ij}^R$  of the matrix, system is written as follows:

$$\begin{aligned} x_{11}v_{11}^R + u_{11}^R z_{11} + x_{12}v_{21}^R + u_{12}^R z_{21} + x_{13}v_{31}^R + u_{13}^R z_{31} &= \delta, \\ x_{11}v_{12}^R + u_{11}^R z_{12} + x_{12}v_{22}^R + u_{12}^R z_{22} + x_{13}v_{32}^R + u_{13}^R z_{32} &= \alpha, \\ x_{11}v_{13}^R + u_{11}^R z_{13} + x_{12}v_{23}^R + u_{12}^R z_{23} + x_{13}v_{33}^R + u_{13}^R z_{33} &= \alpha, \\ \\ x_{21}v_{11}^R + u_{21}^R z_{11} + x_{22}v_{21}^R + u_{22}^R z_{21} + x_{23}v_{31}^R + u_{23}^R z_{31} &= \alpha, \\ x_{21}v_{12}^R + u_{21}^R z_{12} + x_{22}v_{22}^R + u_{22}^R z_{22} + x_{23}v_{32}^R + u_{23}^R z_{32} &= \delta, \\ x_{21}v_{13}^R + u_{21}^R z_{13} + x_{22}v_{23}^R + u_{22}^R z_{23} + x_{23}v_{33}^R + u_{23}^R z_{33} &= \alpha, \\ \\ x_{31}v_{11}^R + u_{31}^R z_{11} + x_{32}v_{21}^R + u_{32}^R z_{21} + x_{33}v_{31}^R + u_{33}^R z_{31} &= \alpha, \\ x_{31}v_{12}^R + u_{31}^R z_{12} + x_{32}v_{22}^R + u_{32}^R z_{22} + x_{33}v_{32}^R + u_{33}^R z_{32} &= \alpha, \\ x_{31}v_{13}^R + u_{31}^R z_{13} + x_{32}v_{23}^R + u_{32}^R z_{23} + x_{33}v_{33}^R + u_{33}^R z_{33} &= \delta. \end{aligned}$$

**Step 4:**

Form the matrix  $\tilde{Z}$  by substituting the values of  $z_{ij}$ ,  $v_{ij}^L$  and  $v_{ij}^R$ .

**Remark 1:**

If only center parts

$$X = \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix} \text{ and } Z = \begin{pmatrix} z_{11} & z_{12} & z_{13} \\ z_{21} & z_{22} & z_{23} \\ z_{31} & z_{32} & z_{33} \end{pmatrix}$$

of  $\tilde{X}$  and  $\tilde{Z}$  respectively are considered, then  $XZ = I$  or  $ZX = I$  i.e.  $Z$  is inverse of  $X$ .

**Remark 2:**

The final opinion obtained in this way is not unique since the uniqueness of fuzzy inverse matrix is not guaranteed.

**Example**

Consider the initial opinion of 3 experts:

$$\tilde{y}^{(1)} = \begin{pmatrix} (0,5,1,2) \\ (1,3,4) \\ (2,3,5) \end{pmatrix}$$

about some alternative, and

$$\tilde{W} = \begin{pmatrix} (1,3,5) & (2,4,5) & (0,2,3) \\ (0,1,2) & (2,4,6) & (3,4,7) \\ (1,2,5) & (3,4,6) & (0,2,5) \end{pmatrix}$$

be the matrix of interpersonal influences of these experts. Computation of final opinions is required.

**Solution**

**Step 1:** Normalized matrix of interpersonal influences is obtained by using operations defined in section 2.1.1 (see appendix):

$$\tilde{W}^{(N)} = \begin{pmatrix} \left(\frac{1}{13}, \frac{1}{3}, \frac{5}{3}\right) & \left(\frac{2}{13}, \frac{4}{9}, \frac{5}{3}\right) & \left(0, \frac{2}{9}, 1\right) \\ \left(0, \frac{1}{9}, \frac{2}{5}\right) & \left(\frac{2}{15}, \frac{4}{9}, \frac{6}{5}\right) & \left(\frac{1}{5}, \frac{4}{9}, \frac{7}{5}\right) \\ \left(\frac{1}{16}, \frac{1}{4}, \frac{5}{4}\right) & \left(\frac{3}{16}, \frac{1}{2}, \frac{3}{2}\right) & \left(0, \frac{1}{4}, \frac{5}{4}\right) \end{pmatrix}$$

$$\tilde{A} = \begin{pmatrix} \left(\frac{2}{13}, \frac{2}{3}, \frac{8}{3}\right) & (0,0,0) & (0,0,0) \\ (0,0,0) & \left(\frac{1}{5}, \frac{5}{9}, \frac{9}{5}\right) & (0,0,0) \\ (0,0,0) & (0,0,0) & \left(\frac{1}{4}, \frac{3}{4}, \frac{11}{4}\right) \end{pmatrix}$$

In this step identity matrix is considered while in the

next step, fuzzy identity matrix is considered.

**Step 2:**

$$\tilde{A}\tilde{W} = \begin{pmatrix} \left(\frac{2}{169}, \frac{2}{9}, \frac{40}{9}\right) & \left(\frac{4}{169}, \frac{8}{27}, \frac{40}{9}\right) & \left(0, \frac{4}{27}, \frac{8}{3}\right) \\ \left(0, \frac{5}{81}, \frac{18}{25}\right) & \left(\frac{2}{75}, \frac{20}{81}, \frac{62}{25}\right) & \left(\frac{1}{25}, \frac{20}{81}, \frac{62}{25}\right) \\ \left(\frac{1}{64}, \frac{3}{16}, \frac{55}{16}\right) & \left(\frac{3}{64}, \frac{3}{8}, \frac{33}{8}\right) & \left(0, \frac{3}{16}, \frac{55}{16}\right) \end{pmatrix}$$

$$(\tilde{I} - \tilde{A}) = \begin{pmatrix} (1 - \delta, 1, 1 + \delta) & (-\alpha, 0, \alpha) & (-\alpha, 0, \alpha) \\ (-\alpha, 0, \alpha) & (1 - \delta, 1, 1 + \delta) & (-\alpha, 0, \alpha) \\ (-\alpha, 0, \alpha) & (-\alpha, 0, \alpha) & (1 - \delta, 1, 1 + \delta) \end{pmatrix}$$

$$= \begin{pmatrix} \left(\frac{2}{13}, \frac{2}{3}, \frac{8}{3}\right) & (0,0,0) & (0,0,0) \\ (0,0,0) & \left(\frac{1}{5}, \frac{5}{9}, \frac{9}{5}\right) & (0,0,0) \\ (0,0,0) & (0,0,0) & \left(\frac{1}{4}, \frac{3}{4}, \frac{11}{4}\right) \end{pmatrix}$$

$$= \begin{pmatrix} \left(\frac{11}{13} - \delta, \frac{1}{3}, \delta - \frac{5}{3}\right) & (-\alpha, 0, \alpha) & (-\alpha, 0, \alpha) \\ (-\alpha, 0, \alpha) & \left(\frac{4}{5} - \delta, \frac{4}{9}, \delta - \frac{4}{5}\right) & (-\alpha, 0, \alpha) \\ (-\alpha, 0, \alpha) & (-\alpha, 0, \alpha) & \left(\frac{3}{4} - \delta, \frac{1}{4}, \delta - \frac{7}{4}\right) \end{pmatrix}$$

$$\tilde{I} - \tilde{A}\tilde{W} = \begin{pmatrix} \left(\frac{167}{169} - \delta, \frac{7}{9}, \delta - \frac{31}{9}\right) & \left(-\alpha - \frac{4}{169}, \frac{-8}{27}, \alpha - \frac{40}{9}\right) & \left(-\alpha, \frac{-4}{27}, \alpha - \frac{8}{3}\right) \\ \left(-\alpha, \frac{-5}{81}, \alpha - \frac{18}{25}\right) & \left(\frac{73}{75} - \delta, \frac{61}{81}, \delta - \frac{38}{25}\right) & \left(-\alpha - \frac{1}{25}, \frac{-20}{81}, \alpha - \frac{63}{25}\right) \\ \left(-\alpha - \frac{1}{64}, \frac{-3}{16}, \alpha - \frac{55}{16}\right) & \left(-\alpha - \frac{3}{64}, \frac{-3}{8}, \alpha - \frac{33}{8}\right) & \left(1 - \delta, \frac{13}{16}, \delta - \frac{39}{16}\right) \end{pmatrix}$$

**Step 3:**

$$\text{Let } \tilde{Z} = \begin{pmatrix} (z_{11} - v_{11}^L, z_{11}, z_{11} + v_{11}^R) & (z_{12} - v_{12}^L, z_{12}, z_{12} + v_{12}^R) & (z_{13} - v_{13}^L, z_{13}, z_{13} + v_{13}^R) \\ (z_{21} - v_{21}^L, z_{21}, z_{21} + v_{21}^R) & (z_{22} - v_{22}^L, z_{22}, z_{22} + v_{22}^R) & (z_{23} - v_{23}^L, z_{23}, z_{23} + v_{23}^R) \\ (z_{31} - v_{31}^L, z_{31}, z_{31} + v_{31}^R) & (z_{32} - v_{32}^L, z_{32}, z_{32} + v_{32}^R) & (z_{33} - v_{33}^L, z_{33}, z_{33} + v_{33}^R) \end{pmatrix}$$

be the inverse of  $\tilde{I} - \tilde{A}\tilde{W}$ , described above. Then to find the

center part of the matrix, following system of equations is solved:

$$\begin{aligned} \frac{7}{9}z_{11} - \frac{8}{27}z_{21} - \frac{4}{27}z_{31} &= 1, \\ \frac{5}{81}z_{11} + \frac{61}{81}z_{21} - \frac{20}{81}z_{31} &= 0, \\ \frac{3}{16}z_{11} - \frac{3}{8}z_{21} + \frac{13}{16}z_{31} &= 0, \end{aligned}$$

$$\begin{aligned} \frac{7}{9}z_{12} - \frac{8}{27}z_{22} - \frac{4}{27}z_{32} &= 0, \\ \frac{5}{81}z_{12} + \frac{61}{81}z_{22} - \frac{20}{81}z_{32} &= 1, \\ \frac{3}{16}z_{12} - \frac{3}{8}z_{22} + \frac{13}{16}z_{32} &= 0, \end{aligned}$$

$$\begin{aligned} \frac{7}{9}z_{13} - \frac{8}{27}z_{23} - \frac{4}{27}z_{33} &= 0, \\ \frac{5}{81}z_{13} + \frac{61}{81}z_{23} - \frac{20}{81}z_{33} &= 0, \\ \frac{3}{16}z_{13} - \frac{3}{8}z_{23} + \frac{13}{16}z_{33} &= 1, \end{aligned}$$

$$\begin{aligned} z_{11} &= 1.4796, z_{12} = 0.8442, z_{13} = 0.5263, \\ z_{21} &= 0.2748, z_{22} = 1.7214, z_{23} = 0.5732, \\ z_{31} &= 0.4682, z_{32} = 0.9893, z_{33} = 1.6168. \end{aligned}$$

To find the left spread part, following system of equations is solved:

$$\begin{aligned} \frac{7}{9}v_{11}^L + \left(\frac{167}{169} - \delta\right)z_{11} - \frac{8}{27}v_{21}^L + \left(-\frac{4}{169} - \alpha\right)z_{21} - \frac{4}{27}v_{31}^L + (-\alpha)z_{31} &= \delta, \\ \frac{5}{81}v_{11}^L + (-\alpha)z_{11} - \frac{61}{81}v_{21}^L + \left(\frac{73}{75} - \delta\right)z_{21} - \frac{20}{81}v_{31}^L + \left(-\frac{1}{25} - \alpha\right)z_{31} &= \alpha, \\ \frac{3}{16}v_{11}^L + \left(-\frac{1}{64} - \alpha\right)z_{11} - \frac{3}{8}v_{21}^L + \left(-\frac{3}{64} - \alpha\right)z_{21} + \frac{13}{16}v_{31}^L + (1 - \delta)z_{31} &= \alpha, \end{aligned}$$

$$\begin{aligned} \frac{7}{9}v_{12}^L + \left(\frac{167}{169} - \delta\right)z_{12} - \frac{8}{27}v_{22}^L + \left(-\frac{4}{169} - \alpha\right)z_{22} - \frac{4}{27}v_{32}^L + (-\alpha)z_{32} &= \alpha, \\ \frac{5}{81}v_{12}^L + (-\alpha)z_{12} - \frac{61}{81}v_{22}^L + \left(\frac{73}{75} - \delta\right)z_{22} - \frac{20}{81}v_{32}^L + \left(-\frac{1}{25} - \alpha\right)z_{32} &= \delta, \\ \frac{3}{16}v_{12}^L + \left(-\frac{1}{64} - \alpha\right)z_{12} - \frac{3}{8}v_{22}^L + \left(-\frac{3}{64} - \alpha\right)z_{22} + \frac{13}{16}v_{32}^L + (1 - \delta)z_{32} &= \alpha, \end{aligned}$$

$$\begin{aligned} \frac{7}{9}v_{13}^L + \left(\frac{167}{169} - \delta\right)z_{13} - \frac{8}{27}v_{23}^L + \left(-\frac{4}{169} - \alpha\right)z_{23} - \frac{4}{27}v_{33}^L + (-\alpha)z_{33} &= \alpha, \\ \frac{5}{81}v_{13}^L + (-\alpha)z_{13} - \frac{61}{81}v_{23}^L + \left(\frac{73}{75} - \delta\right)z_{23} - \frac{20}{81}v_{33}^L + \left(-\frac{1}{25} - \alpha\right)z_{33} &= \alpha, \\ \frac{3}{16}v_{13}^L + \left(-\frac{1}{64} - \alpha\right)z_{13} - \frac{3}{8}v_{23}^L + \left(-\frac{3}{64} - \alpha\right)z_{23} + \frac{13}{16}v_{33}^L + (1 - \delta)z_{33} &= \delta. \end{aligned}$$

To find the right spread part, following system of equations is solved:

$$\begin{aligned} \frac{7}{9}v_{11}^R + \left(\delta - \frac{31}{9}\right)z_{11} - \frac{8}{27}v_{21}^R + \left(\alpha - \frac{40}{9}\right)z_{21} - \frac{4}{27}v_{31}^R + \left(\alpha - \frac{8}{3}\right)z_{31} &= \delta, \\ \frac{5}{81}v_{11}^R + \left(\alpha - \frac{18}{25}\right)z_{11} - \frac{61}{81}v_{21}^R + \left(\delta - \frac{38}{25}\right)z_{21} - \frac{20}{81}v_{31}^R + \left(\alpha - \frac{63}{25}\right)z_{31} &= \alpha, \\ \frac{3}{16}v_{11}^R + \left(\alpha - \frac{55}{16}\right)z_{11} - \frac{3}{8}v_{21}^R + \left(\alpha - \frac{33}{8}\right)z_{21} + \frac{13}{16}v_{31}^R + \left(\delta - \frac{39}{16}\right)z_{31} &= \alpha, \end{aligned}$$

$$\begin{aligned} \frac{7}{9}v_{12}^R + \left(\delta - \frac{31}{9}\right)z_{12} - \frac{8}{27}v_{22}^R + \left(\alpha - \frac{40}{9}\right)z_{22} - \frac{4}{27}v_{32}^R + \left(\alpha - \frac{8}{3}\right)z_{32} &= \alpha, \\ \frac{5}{81}v_{12}^R + \left(\alpha - \frac{18}{25}\right)z_{12} - \frac{61}{81}v_{22}^R + \left(\delta - \frac{38}{25}\right)z_{22} - \frac{20}{81}v_{32}^R + \left(\alpha - \frac{63}{25}\right)z_{32} &= \delta, \\ \frac{3}{16}v_{12}^R + \left(\alpha - \frac{55}{16}\right)z_{12} - \frac{3}{8}v_{22}^R + \left(\alpha - \frac{33}{8}\right)z_{22} + \frac{13}{16}v_{32}^R + \left(\delta - \frac{39}{16}\right)z_{32} &= \alpha, \end{aligned}$$

$$\begin{aligned} \frac{7}{9}v_{13}^R + \left(\delta - \frac{31}{9}\right)z_{13} - \frac{8}{27}v_{23}^R + \left(\alpha - \frac{40}{9}\right)z_{23} - \frac{4}{27}v_{33}^R + \left(\alpha - \frac{8}{3}\right)z_{33} &= \alpha, \\ \frac{5}{81}v_{13}^R + \left(\alpha - \frac{18}{25}\right)z_{13} - \frac{61}{81}v_{23}^R + \left(\delta - \frac{38}{25}\right)z_{23} - \frac{20}{81}v_{33}^R + \left(\alpha - \frac{63}{25}\right)z_{33} &= \alpha, \\ \frac{3}{16}v_{13}^R + \left(\alpha - \frac{55}{16}\right)z_{13} - \frac{3}{8}v_{23}^R + \left(\alpha - \frac{33}{8}\right)z_{23} + \frac{13}{16}v_{33}^R + \left(\delta - \frac{39}{16}\right)z_{33} &= \delta. \end{aligned}$$

Finally,

$$(\tilde{I} - \tilde{A}\tilde{W})^{-1} = \begin{pmatrix} (4.0708 - 4.14736 - 5.0379\alpha), & (3.8706 - 4.06736 - 8.9151\alpha), & (2.5247 - 2.64006 - 7.9523\alpha), \\ 1.4796, & 0.8442, & 0.5247, \\ (1.4813 - 2.2968\alpha - 1.1881\delta), & (31.7316 - 4.9032\alpha - 2.3789\delta), & (22.1241 - 3.3046\alpha - 1.5873\delta), \\ (1.3508 - 1.42296 - 0.6857\alpha), & (5.2689 - 5.48396 - 6.2201\alpha), & (2.4682 - 2.63156 - 6.9177\alpha), \\ 0.2748, & 1.7214, & 0.5732, \\ (11.1583 - 2.2694\alpha - 0.8733\delta), & (22.307 - 4.5240\alpha - 2.0410\delta), & (16.7743 - 2.9252\alpha - 1.4850\delta), \\ (2.0948 - 2.1901\delta - 7.7177\alpha), & (4.4271 - 4.68736 - 9.3167\alpha), & (4.8993 - 5.04446 - 6.3812\alpha), \\ 0.4682, & 0.9893, & 1.6168, \\ (18.5478 - 2.5055\alpha - 1.2535\delta), & (32.8976 - 5.1465\alpha - 2.7086\delta), & (24.0659 - 3.4660\alpha - 1.8108\delta) \end{pmatrix}$$

$$(\tilde{I} - \tilde{A}\tilde{W})^{-1}(\tilde{I} - \tilde{A}) = \begin{pmatrix} (4.0708 - 4.14736 - 5.0379\alpha), & (3.8706 - 4.06736 - 8.9151\alpha), & (2.5247 - 2.646 - 7.9523\alpha), \\ (0.8461 - \delta) - \alpha, & (0.8 - \delta) - \alpha, & (0.75 - \delta) - \alpha, \\ (6.3953 - 6.70736 - 18.6874\alpha), & (6.5955 - 6.78736 - 12.9902\alpha), & (7.9414 - 8.21466 - 13.953\alpha), \\ \frac{1}{3}(1.4796), & \frac{4}{9}(0.8442), & \frac{1}{4}(0.5263), \\ (1.4813 - 2.2968\alpha - 1.1881\delta), & (31.7316 - 4.9032\alpha - 2.3789\delta), & (22.1241 - 3.3046\alpha - 1.5873\delta), \\ (\delta - 1.6666) + \alpha, & (\delta - 0.8) + \alpha, & (\delta - 1.75) + \alpha, \\ (53.8557 - 8.2078\alpha - 3.9662\delta), & (23.6054 - 5.6014\alpha - 2.7754\delta), & (33.2129 - 7.2\alpha - 3.567\delta), \\ (1.3508 - 1.42296 - 0.6857\alpha), & (5.2689 - 6.2201\alpha - 5.4839\delta), & (2.4682 - 2.63156 - 6.9177\alpha), \\ (0.8461 - \delta) - \alpha, & (0.8 - \delta) - \alpha, & (0.75 - \delta) - \alpha, \\ (7.7371 - 13.1378\alpha - 8.1154\delta), & (3.819 - 4.05446 - 7.6034\alpha), & (6.6197 - 6.90688 - 6.9058\alpha), \\ \frac{1}{3}(0.2748), & \frac{4}{9}(1.7214), & \frac{1}{4}(0.5732), \\ (11.1583 - 2.2694\alpha - 0.8733\delta), & (22.307 - 4.5240\alpha - 2.0410\delta), & (16.7743 - 2.9252\alpha - 1.4850\delta), \\ (\delta - 1.6666) + \alpha, & (\delta - 0.8) + \alpha, & (\delta - 1.75) + \alpha, \\ (39.0813 - 7.4492\alpha - 3.526\delta), & (27.9326 - 5.1936\alpha - 2.3583\delta), & (23.4653 - 6.7924\alpha - 2.914\delta), \\ (2.0948 - 2.1901\delta - 7.7177\alpha), & (4.4271 - 4.68736 - 9.3167\alpha), & (4.8993 - 5.04446 - 6.3812\alpha), \\ (0.8461 - \delta) - \alpha, & (0.8 - \delta) - \alpha, & (0.75 - \delta) - \alpha, \\ (9.3264 - 9.7317\delta - 15.6979\alpha), & (6.9941 - 7.23456 - 14.0989\alpha), & (6.5219 - 6.87746 - 17.0344\alpha), \\ \frac{1}{3}(0.4682), & \frac{4}{9}(0.9893), & \frac{1}{4}(1.6168), \\ (18.5478 - 2.5055\alpha - 1.2535\delta), & (32.8976 - 5.1465\alpha - 2.7086\delta), & (24.0659 - 3.4660\alpha - 1.8181\delta), \\ (\delta - 1.6666) + \alpha, & (\delta - 0.8) + \alpha, & (\delta - 1.75) + \alpha, \\ (56.9635 - 8.6125\alpha - 4.5194\delta), & (42.6137 - 5.9715\alpha - 3.0643\delta), & (51.4454 - 7.652\alpha - 3.9621\delta) \end{pmatrix}$$

$$(\tilde{I} - \tilde{A}\tilde{W})^{-1}(\tilde{I} - \tilde{A})^{(1)} = \begin{pmatrix} 8.6056 - 19.9237\delta - 46.8677\alpha + 53.9088\delta\alpha + 11.4209\delta^2 + 49.3299\alpha^2, \\ -300.0644 + 265.9710\delta + 420.4570\alpha - 77.5984\delta\alpha - 19.8283\delta^2 - 74.8212\alpha^2, \\ 8.4888 - 19.8176\delta - 36.5696\alpha + 42.3240\delta\alpha + 11.4583\delta^2 + 27.9839\alpha^2, \\ 2.8167, \\ -255.3503 + 2178519\delta + 354.852\alpha - 68.314\delta\alpha - 17.3356\delta^2 - 69.6348\alpha^2, \\ 11.7768 - 27.5159\delta - 44.9912\alpha + 51.5904\delta\alpha + 15.8711\delta^2 + 56.0166\alpha^2, \\ 2.6877, \\ -377.6724 + 317.7056\delta + 596.7564\alpha - 87.0978\delta\alpha - 22.3954\delta^2 - 79.371\alpha^2 \end{pmatrix}$$

which is the final opinion of experts.

### CONCLUSION

In this paper, CRP is remodeled with fuzzy numbers, instead of crisp numbers. There are some limitations in the example given for illustration. In step 1, diagonal entries  $\tilde{a}_{ii}$ ,  $i = 1, 2, 3$  of the matrix  $\tilde{A}$  are equal to  $\tilde{1} - \tilde{w}_{ii}$ .  $\tilde{1}$  is chosen according to the entries of  $\tilde{W}^{(N)}$  i.e.  $(\frac{3}{13}, 1, \frac{13}{3}), (\frac{1}{5}, 1, 3), (\frac{1}{4}, 1, 4)$  are the respective fuzzy one numbers for three rows. In step 2, fuzzy identity matrix  $\tilde{I}$  consists of three equal fuzzy one numbers and six equal fuzzy zero numbers. This matrix can also be chosen with different fuzzy one and fuzzy zero numbers (means  $\delta$  and  $\alpha$  are not necessarily same for each entry). As well as decision-making problems are capable to model with fuzzy numbers, interpersonal influences and opinion dynamics have also the potential to process with fuzzy numbers and fuzzy matrices in more general forms. Moreover some other types and generalizations of fuzzy numbers would contribute to the proceedings of influences and opinions in future. This work can facilitate some other consensus-reaching processes with fuzzy numbers and prepare for advancements in fuzzy inverse matrices.

### AUTHORSHIP CONTRIBUTIONS

Authors equally contributed to this work.

### DATA AVAILABILITY STATEMENT

The authors confirm that the data that supports the findings of this study are available within the article. Raw

data that support the finding of this study are available from the corresponding author, upon reasonable request.

## CONFLICT OF INTEREST

The author declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## ETHICS

There are no ethical issues with the publication of this manuscript.

## REFERENCES

- [1] Palomares I, Estrella FJ, Martínez L, Herrera F. Consensus under a fuzzy context: taxonomy, analysis framework AFRYCA and experimental case of study. *Inf Fusion* 2014;20:252–271. [\[CrossRef\]](#)
- [2] Herrera-Viedma E, Cabrerizo FJ, Kacprzyk J, Pedrycz W. A review of soft consensus models in a fuzzy environment. *Inf Fusion* 2014;17:4–13. [\[CrossRef\]](#)
- [3] Dong Y, Zha Q, Zhang H, Kou G, Fujita H, Chiclana F, et al. Consensus reaching in social network group decision making: research paradigms and challenges. *Knowl Based Syst* 2018;162:3–13. [\[CrossRef\]](#)
- [4] Su W, Zhang L, Zeng S, Jin H. A fuzzy-social network multi-criteria group decision-making framework for selection of renewable energy project: a case of China. *Int J Fuzzy Syst* 2021;24:1059–1078. [\[CrossRef\]](#)
- [5] Capuano N, Chiclana F, Fujita H, Herrera-Viedma E, Loia V. Fuzzy group decision making with incomplete information guided by social influence. *IEEE Trans Fuzzy Syst* 2017;26:1704–1718. [\[CrossRef\]](#)
- [6] Yao S, Gu M. An influence network-based consensus model for large-scale group decision making with linguistic information. *Int J Comput Intell Syst* 2022;15:3. [\[CrossRef\]](#)
- [7] Friedkin N, Johnson E. Social influence network and opinion change. *Adv Group Process* 1999;16:1–29.
- [8] Proskurnikov AV, Tempo R, Cao M, Friedkin NE. Opinion evolution in time-varying social influence networks with prejudiced agents. *IFAC Pap Online* 2017;50:11896–11901. [\[CrossRef\]](#)
- [9] Perez LG, Mata F, Chiclana F, Kou G, Herrera-Viedma E. Modelling influence in group decision making. *Soft Comput* 2016;20:1653–1665. [\[CrossRef\]](#)
- [10] Tsuji R. Interpersonal influence and attitude change toward conformity in small groups: a social psychological model. *J Math Sociol* 2010;17–34. [\[CrossRef\]](#)
- [11] Dong Y, Zhan M, Kou G, Ding Z, Liang H. A survey on the fusion process in opinion dynamics. *Inf Fusion* 2018;43:57–65. [\[CrossRef\]](#)
- [12] Zadeh LA. Fuzzy sets. *Inf Control* 1965;8:338–353. [\[CrossRef\]](#)
- [13] Mohanta KK, Sharanappa DS, Dabke D, Mishra LN, Mishra VN. Data envelopment analysis in the context of spherical fuzzy inputs and outputs. *Eur J Pure Appl Math* 2022;15:1158–1179. [\[CrossRef\]](#)
- [14] Sharma MK, Sadhna, Bhargava AK, Kumar S, Rathour L, Mishra LN, Pandey S. A Fermatean fuzzy ranking function in optimization of intuitionistic fuzzy transportation problems. *Adv Math Models Appl* 2022;7:191–204.
- [15] Pakhira R, Ghosh U, Sarkar S, Mishra LN. Study of memory effect in an EOQ model with fractional polynomial demand rate under fuzzy environment. *Discontinuity Nonlinearity Complex* 2022;11:583–598. [\[CrossRef\]](#)
- [16] Khalid A, Beg I. Influence model of evasive decision makers. *J Intell Fuzzy Syst* 2019;37:2539–2548. [\[CrossRef\]](#)
- [17] Mahmood A, Abbas M. Influence model and doubly extended TOPSIS with TOPSIS-based matrix of interpersonal influences. *J Intell Fuzzy Syst* 2020;39:7537–7546. [\[CrossRef\]](#)
- [18] Mahmood A, Abbas M. Influence model with opinions and trust score evaluations under the leader-follower environment. *J Intell Fuzzy Syst* 2021;41:6363–6373. [\[CrossRef\]](#)
- [19] Jain R. A procedure for multi-aspect decision making using fuzzy sets. *Int J Syst Sci* 1978;8:1–7. [\[CrossRef\]](#)
- [20] Dubois D, Prade H. Operations on fuzzy numbers. *Int J Syst Sci* 1978;9:613–626. [\[CrossRef\]](#)
- [21] Abasbandy S, Asady B. Ranking of fuzzy numbers by sign distance. *Inf Sci* 2006;176:2405–2416. [\[CrossRef\]](#)
- [22] Bortolan G, Degani R. A review of some methods for ranking fuzzy numbers. *Fuzzy Sets Syst* 1985;15:1–19. [\[CrossRef\]](#)
- [23] Cheng CH. A new approach for ranking fuzzy numbers by distance method. *Fuzzy Sets Syst* 1998;95:307–317. [\[CrossRef\]](#)
- [24] Chu TC, Tsao CT. Ranking fuzzy numbers with an area between the centroid point and original point. *Comput Math Appl* 2002;43:111–117. [\[CrossRef\]](#)
- [25] Ross TJ. Fuzzy arithmetic and the extension principle. In: *Fuzzy logic with engineering applications*. 3rd ed. Chichester: John Wiley & Sons; 2010.
- [26] Lee KH. Fuzzy number. In: *First course on fuzzy theory and applications*. Berlin: Springer; 2005. p. 129–151.
- [27] Amin F, Fahmi A, Abdullah S. Dealer using a new trapezoidal cubic hesitant fuzzy TOPSIS method and application to group decision-making program. *Soft Comput* 2018;23:5353–5366. [\[CrossRef\]](#)
- [28] Akram M, Arshad M. A novel trapezoidal bipolar fuzzy TOPSIS method for group decision-making. *Group Decis Negot* 2019;28:565–584. [\[CrossRef\]](#)
- [29] Rohn J. Interval matrices: singularity and eigenvalues. *SIAM J Matrix Anal Appl* 1993;14:82–91. [\[CrossRef\]](#)
- [30] Rohn J. Checking properties of interval matrices. *Tech Rep* 1996;686.

- [31] Karthick P, Mohiuddine SA, Tamilvanan K, Narayanamoorthy S, Maheswan S. Investigations of color image segmentation based on connectivity measure, shape priority and normalized fuzzy graph cut. *Appl Soft Comput* 2023;139:110239. [\[CrossRef\]](#)
- [32] Gustafson DE, Kessel WC. Fuzzy clustering with a fuzzy covariance matrix. *Proc IEEE*; 1978. p. 761–766. [\[CrossRef\]](#)
- [33] Sharma MK, Dhiman N, Mishra VN, Mishra LN, Dhaka A, Koundal D. Post-symptomatic detection of COVID-2019 grade-based mediative fuzzy projection. *Comput Electr Eng J* 2022;101:108028. [\[CrossRef\]](#)
- [34] Sharma MK, Dhiman N, Mishra LN, Mishra VN, Sahani SK. Mediative fuzzy extension technique and its consistent measurement in the decision making of medical application. *Math Probl Eng* 2021;2021:5530681. [\[CrossRef\]](#)
- [35] Ghaoui LE. Inversion error, condition number, and approximate inverses of uncertain matrices. *Linear Algebra Appl* 2002;343-4:171–193. [\[CrossRef\]](#)
- [36] Dehghan M, Ghatee M, Hashemi B. Inverse of a fuzzy matrix of fuzzy numbers. *Int J Comput Math* 2009;86:1433–1452. [\[CrossRef\]](#)
- [37] Farahani H, Ebadi MJ, Jafari H. Finding inverse of a fuzzy matrix using eigenvalue method. *Int J Innov Technol Explor Eng* 2019;9:3030–3037. [\[CrossRef\]](#)
- [38] Dequan S, Guo X. Solving fuzzy linear matrix equation. *J Phys Conf Ser* 2020;1592:012051. [\[CrossRef\]](#)
- [39] Anandhkumar M, Punithavalli G, Soupramanien T, Broumi S. Generalized symmetric neutrosophic fuzzy matrices. *Neutrosophic Sets Syst* 2023;57:6.
- [40] Anandhkumar M, Harikrishnan T, Chithra SM, Kamalakannan V, Kanimozhi B, Broumi S, et al. Reverse Sharp and Left-T Right-T partial ordering on neutrosophic fuzzy matrices. *Int J Neutrosophic Sci* 2023;21:135–145.
- [41] Basaran MA. Calculating fuzzy inverse matrix using fuzzy linear equation system. *Appl Soft Comput* 2012;12:1810–1813. [\[CrossRef\]](#)
- [42] Tang HC. Decomposition and intersection of two fuzzy numbers for fuzzy preference relations. *Symmetry* 2017;228:9. [\[CrossRef\]](#)
- [43] Wang W, Wang Z. Total orderings defined on the set of all fuzzy numbers. *Fuzzy Sets Syst* 2014;243:131–141. [\[CrossRef\]](#)
- [44] Dehghan M, Hashemi B, Ghatee M. Computational methods for solving fully fuzzy linear systems. *Appl Math Comput* 2006;179:328–343. [\[CrossRef\]](#)

## APPENDIX

Consider the first row of  $\tilde{W}$ :

$$\begin{aligned}
 (1,3,5) \oplus (2,4,5) \oplus (0,2,3) &= (3,9,13) \\
 (1,3,5)/(3,9,13) &= (1/13,1/3,5/3) \\
 (2,4,5)/(3,9,13) &= (2/13,4/9,5/3) \\
 (0,2,3)/(3,9,13) &= (0,2/9,1)
 \end{aligned}$$

And  $(1/13,1/3,5/3) \oplus (2/13,4/9,5/3) \oplus (0,2/9,1) = (3/13,1,13/3)$ , which is a fuzzy one number as defined in Definition 3.